

Noncommutative renormalization for massless QED

Christian Brouder

Laboratoire de Minéralogie-Cristallographie, CNRS UMR7590, Universités Paris 6 et 7,
IPGP, 4 place Jussieu, 75252 Paris Cedex 05, France. brouder@lmcp.jussieu.fr

Alessandra Frabetti

Institut de Mathématique, Bâtiment de Chimie, Université de Lausanne,
CH-1015 Lausanne, Switzerland afrabett@magma.unil.ch

February 1, 2008

Abstract

We study the renormalization of massless QED from the point of view of the Hopf algebra discovered by D. Kreimer. For QED, we describe a Hopf algebra of renormalization which is neither commutative nor cocommutative. We obtain explicit renormalization formulas for the electron and photon propagators, for the vacuum polarization and the electron self-energy, which are equivalent to Zimmermann's forest formula for the sum of all Feynman diagrams at a given order of interaction.

Then we extend to QED the Connes-Kreimer map defined by the coupling constant of the theory (i.e. the homomorphism between some formal diffeomorphisms and the Hopf algebra of renormalization) by defining a noncommutative Hopf algebra of diffeomorphisms, and then showing that the renormalization of the electric charge defines a homomorphism between this Hopf algebra and the Hopf algebra of renormalization of QED.

Finally we show that Dyson's formulas for the renormalization of the electron and photon propagators can be given in a noncommutative (e.g. matrix-valued) form.

Introduction

D. Kreimer made the remarkable discovery that the combinatorics of renormalization was hiding a Hopf algebra [19]. The publication of ref.[20] has spurred many developments in the theory of renormalization and in the application of Hopf algebras to quantum field theories. Among these developments, one of the most intriguing is the construction, by A. Connes and D. Kreimer, of a homomorphism between the Hopf algebra of renormalization of the massless φ^3 quantum field theory and the Connes-Moscovici Hopf algebra of some formal diffeomorphisms on the complex line (see [12]). Such a homomorphism will be called a Connes-Kreimer map. In the present paper, we build a Connes-Kreimer map for the case of the noncommutative (nor cocommutative) Hopf algebra of renormalization of massless quantum electrodynamics (QED).

In the φ^3 quantum field theory, the field φ is scalar. Thus, the amplitudes given by Feynman diagrams are scalars, and their product is commutative. In QED, the fields are either spinors (for the electrons) or vectors (for the photons), and the amplitudes are 4×4 complex matrices. Therefore, their product is not commutative. If we want the Hopf algebra describing the renormalization of QED to be compatible with the product of amplitudes, the algebraic product cannot be commutative. This choice was made in Ref.[7], where we described the renormalization of massless QED by means of a coproduct on planar binary trees.

In the present paper, we first show that this coproduct defines a noncommutative nor cocommutative Hopf algebra over trees. Then we define the natural noncommutative extension of the Connes-Moscovici algebra of formal diffeomorphisms and we prove that this is also a Hopf algebra. Finally, we prove that the relation between the bare and renormalized electric charge $e_0 = e/\sqrt{Z_3(e)}$ defines a homomorphism between the noncommutative Connes-Moscovici algebra and the Hopf algebra of renormalization of QED.

By “noncommutative renormalization” we mean that the noncommutativity of the product of amplitudes is respected by the renormalization method. In practice, the renormalization factors Z multiply the various terms of the Lagrangian, which is a scalar quantity. Thus, all Z are scalars, therefore commutative, quantities. However, the investigation of renormalization by noncommutative multiplication factors has a practical advantage. When various fields are present, renormalization proceeds through the use of various Z_{ij} coupling the different fields. If we group the fields together in a single field vector, then $\{Z_{ij}\}$ becomes a matrix. Noncommutative renormalization can also be used to investigate the transmutation of leptons due to the rotation of the mass matrix [3]. Such a matrix-valued renormalization factor is used even for QED, in the Plymouth approach, to take care of the infrared divergences, cf. [1, 2].

From a mathematical point of view, a morphism between two noncommutative Hopf algebras is richer than the morphism between their abelianizations. It is surprising that such a morphism between the noncommutative Hopf algebra of formal diffeomorphisms and the algebra of renormalization exists, since it does not seem to be required by the standard renormalization methods. Even more surprising is the fact that such a noncommutative version of the Hopf algebra of formal diffeomorphisms exists. Intuitively, we would like to consider it as the algebra of nonscalar functions on formal series with noncommutative coefficients, but this point of view fails for two main reasons: on one side the composition of such series is not associative, while the dual coproduct is, on the other side, even if the composition were associative, the convolution does not preserve the noncommutative characters of the Hopf algebra, and by duality we could not recover the formal series from the Hopf algebra of functions. If we force the composition of series to be associative, by introducing a suitable normal ordering, we come to another surprising fact: the composition in normal ordering is not a binary operation, while the coproduct on the Hopf algebra is. In conclusion, the only way to regard this Hopf algebra is as a deformation of the set of scalar functions on formal series, a kind of infinite dimensional “quantum group” with no commutation relations at all.

The present paper follows the concepts developed in the recent works by Connes and Kreimer. However, since our Hopf algebras are neither commutative nor cocommutative, we cannot use the Milnor-Moore theorem to deal with the associated Lie algebras. Our proofs are elementary, mainly based on recurrence relations and generating functions. They provide explicit expressions for the coproducts and reveal an interesting connection with the resolvent of linear operators.

We do not consider infrared divergences.

Notation. We suppose that all vector spaces and algebras are defined over the field $\mathbb{C}$ of complex numbers. We denote by $\mathbb{C}\langle X \rangle$ the algebra of noncommutative polynomials on the set of variables X , and by $\mathbb{C}[X]$ the algebra of commutative polynomials. The ideal generated by x is denoted by (x) .

When confusion may arise, non-commutative variables are chosen with boldface characters.

Contents

Introduction	1
1 Renormalization “à la Dyson”	3
2 From Feynman graphs to planar binary trees	7
2.1 Correspondence between trees and graphs	8
2.2 Algebra of photon trees and vacuum polarization	12
2.3 Algebra of electron trees and self-energy	14
2.4 Gradings and loops	16
3 Hopf algebras on planar binary trees	18
3.1 Hopf algebra for the photon trees	18
3.2 Hopf structure for the electron trees	20
3.3 Tree expansion for renormalized propagators	22

4	Renormalization “at the order of interaction”	23
4.1	Hopf algebra for photons at order n	24
4.2	Hopf structure for electrons at order n	28
4.3	Renormalization at order n	30
5	Noncommutative Hopf algebra of formal diffeomorphism	32
5.1	The Hopf algebra	32
5.2	Coaction on the dual of series	37
5.3	Normal ordering	38
6	Renormalization of the electric charge	39
6.1	Renormalization of e_0	39
6.2	Renormalization of α_0	44
7	Noncommutative Dyson’s formulas	45
7.1	Dyson’s formula for photons	46
7.2	Dyson’s formula for electrons	48
8	Conclusion and perspective	49
	References	49

1 Renormalization “à la Dyson”

Many experiments proved that the density of charge of an electron placed in the vacuum is in fact modified by the presence of this “vacuum” [17]. Following Dirac, we can consider the vacuum of QED as a medium where all negative energy states are occupied. An electromagnetic field polarizes these negative energy electrons: this is called vacuum polarization, and an electron interacts with its electromagnetic field: this is called electron self-energy [17].

The vacuum polarization and the electron self-energy are usually denoted by Π and Σ , respectively. They can be deduced directly from the Lagrangian through the one-particle Green functions, or propagators, of quantum electrodynamics.

The electron and photon propagators represent the amplitude of probability that the corresponding particle propagates from a point x to a point y , and they are denoted by $S(x, y)$ and $D_{\lambda\mu}(x, y)$ respectively. In vacuum, the propagators are translation-invariant and can be written as $S(x - y)$ and $D_{\lambda\mu}(x - y)$. Their Fourier transforms are denoted by $S(q; e_0)$ and $D(q; e_0)$, where e_0 is the (fictive) electron charge of the electrons “before renormalization” and is called the bare charge.

The vacuum polarization and the self-energy can be deduced from the propagators through the identities

$$D^{-1}(q; e_0) = D_0^{-1}(q; e_0) - \Pi(q; e_0), \quad (1)$$

$$S^{-1}(q; e_0) = S_0^{-1}(q; e_0) - \Sigma(q; e_0), \quad (2)$$

where D^{-1} and S^{-1} are the inverse of the operators D and S .

In fact, many interesting physical properties of the electromagnetic field can be obtained directly from the propagators. From the electron propagator $S(x, y)$, for instance, one can describe the charge density $\rho(x) = i\text{etr}[\gamma^0 S(x, x)]$ and the electromagnetic current $J^k(x) = i\text{etr}[\gamma^k S(x, x)]$. From the photon propagator $D_{\lambda\mu}(x, y)$, instead, one can describe the electromagnetic potential $A_\lambda(x)$ due to an external electromagnetic current $j^\mu(y)$ (to first order) by $A_\lambda(x) = \int dy D_{\lambda\mu}(x, y) j^\mu(y)$. In other words, the photon propagator describes the linear response of the system to an external electromagnetic current.

Most methods available to calculate the propagators are perturbative. In perturbative quantum electrodynamics, the electron self-energy, the vacuum polarization and the propagators are written as

power series over the coupling constant e_0^2 ,

$$\Sigma(q; e_0) = \sum_{n=1}^{\infty} e_0^{2n} \Sigma_n(q), \quad (3)$$

$$\Pi(q; e_0) = \sum_{n=1}^{\infty} e_0^{2n} \Pi_n(q), \quad (4)$$

$$S(q; e_0) = \sum_{n=0}^{\infty} e_0^{2n} S_n(q), \quad (5)$$

$$D(q; e_0) = \sum_{n=0}^{\infty} e_0^{2n} D_n(q). \quad (6)$$

The terms $\Sigma_n(q)$, $\Pi_n(q)$, $S_n(q)$ and $D_n(q)$ are finite sums of integrals. Each of these integrals is classically described by a suitable Feynman diagram Γ with n loops. The graphs Γ should be chosen according to the propagator they describe, and they provide a very figurative and elegant way to remember all possible interactions which should be considered in the expansion at a fixed order. For instance, for $n = 0$ we consider the free propagators, i.e. the propagators for QED without electromagnetic interactions ($e_0 = 0$), which are usually denoted by simple lines

$$S_0(q) = \text{---}, \quad D_0(q) = \text{~~~~~};$$

and for $n = 1$ we consider the graphs with one loop,

$$S_1(q) = \text{---} \text{ (loop) } \text{---}, \quad D_1(q) = \text{~~~~~} \text{ (loop) } \text{~~~~~}.$$

More precisely, denote by $\mathcal{F}^e$ and $\mathcal{F}^\gamma$ the sets of graphs which describe the electron and the photon propagators respectively. The rules to assign an integral to each graph Γ were given by Feynman [14]. Denote by U the ‘‘Feynman rules’’ map, which assign to each graph Γ the suitable integral $U(\Gamma)$. (An explicit description of the sets $\mathcal{F}^e$, $\mathcal{F}^\gamma$ and of the Feynman rules can be found in any textbook of quantum field theory, for instance [17], Appendix 4.) Then, the coefficients $S_n(q)$ and $D_n(q)$ can be described as

$$S_n(q) = \sum_{\Gamma \in \mathcal{F}^e, |\Gamma|=n} U(\Gamma; q),$$

$$D_n(q) = \sum_{\Gamma \in \mathcal{F}^\gamma, |\Gamma|=n} U(\Gamma; q),$$

where $|\Gamma|$ is half the number of vertices in the diagram Γ .

The graphs which contribute to the self-energy and to the vacuum polarization are the so-called 1-particle irreducible (1PI) Feynman diagrams, i.e. those diagrams which can not be seen as junction of two or more meaningful adjacent diagrams. If we denote by $\mathcal{F}_{1PI}^e \subset \mathcal{F}^e$ and $\mathcal{F}_{1PI}^\gamma \subset \mathcal{F}^\gamma$ the subsets of 1PI Feynman diagrams, the coefficients $\Sigma_n(q)$ and $\Pi_n(q)$ can be described as

$$\Sigma_n(q) = \sum_{\Gamma \in \mathcal{F}_{1PI}^e, |\Gamma|=n} U(\Gamma; q),$$

$$\Pi_n(q) = \sum_{\Gamma \in \mathcal{F}_{1PI}^\gamma, |\Gamma|=n} U(\Gamma; q).$$

The problem is that the most of the integrals $U(\Gamma; q)$ involved in these expansions are divergent.

Renormalization theory is a prescription to remove the divergences from the integrals $U(\Gamma; q)$, yielding finite numbers in accordance with experimental data. In practice, renormalization consists in two main steps.

The first step is a regularization of the divergent integrals to avoid manipulating infinities. Many regularizations are possible, the integrals can be limited to an upper value $q^2 < \Lambda^2$ (cutoff), they can be carried out in a non integer dimension $d = 4 - \epsilon$ (dimensional regularization), etc. Various regularizations are described in Refs.[17, 8]. Thus, the terms $U(\Gamma; q)$, and consequently the coefficients $S_n(q)$ and $D_n(q)$ of the perturbative expansions, usually depend on the regularization variables (e.g. Λ , ϵ , etc.) which are left implicit for notational convenience. The original, divergent, integrals are recovered when the regularization parameters tend to the limit which gives the divergency ($\epsilon \rightarrow 0$, $\Lambda \rightarrow \infty$, etc.).

The second step consists in extracting, from the regularized integrals, a term which contains the divergent part, once the regularization parameter is sent to his limit. We take the example of dimensional regularization, for which the regularization parameter is ϵ . To find the finite value corresponding to $U_\epsilon(\Gamma)$, we first remove all subdivergences from $U_\epsilon(\Gamma)$ to obtain a quantity denoted by $\bar{R}_\epsilon(\Gamma)$. In a renormalizable theory, such as QED, the term $\bar{R}_\epsilon(\Gamma)$ has a given order of divergence. For instance, the self-energy is linearly divergent, so that

$$\bar{R}_\epsilon(\Gamma; q) = -C_0(\Gamma, \epsilon) - C_1(\Gamma, \epsilon)q + R_\epsilon(\Gamma; q),$$

where $R_\epsilon(\Gamma; q)$ has now a finite value for $\epsilon \rightarrow 0$, this value being the renormalized value of Γ denoted by $R(\Gamma; q)$. The divergent counterterms $C_0(\Gamma, \epsilon)$ and $C_1(\Gamma, \epsilon)$ are determined by the so-called renormalization conditions, which are boundary conditions for the finite part $R_\epsilon(\Gamma; q)$. For instance, we may require that $R_\epsilon(\Gamma; q)$ and its derivative with respect to q are 0 for $q = 0$. Many other renormalization conditions are possible (cf. [8]). The purpose of the renormalization conditions is determine the finite and the infinite parts of $\bar{R}_\epsilon(\Gamma)$.

For comparison with experiment, the renormalization conditions use experimental results, such as the electron mass or the electron charge [17].

The renormalization of 1-loop diagrams is usually an accessible computation. It becomes a hard task when divergent integrals contain divergent subintegrals, soon as the order of interaction is greater than 1. In such cases, the presence of loops with subloops in Feynman diagrams translates into a dependency of the counterterm of the diagram from those of the subdiagrams.

Renormalization of Feynman diagrams was thoroughly studied, culminating first in Zimmermann's forest formula [32], which describes the intricate dependency of the counterterms of several-loops graphs, and then in Kreimer's discovery of a Hopf algebra structure hidden in the procedure, cf. [19]. Since the renormalization of Feynman diagrams is not our subject, for more details we refer the reader to standard textbooks [17, 8] and to the recent publications [10, 11, 12, 21].

After renormalization, by summing up all the finite integrals we obtain a finite value of the propagators at each order of interaction,

$$\begin{aligned}\bar{S}_n(q) &= \sum_{\Gamma \in \mathcal{F}^\epsilon, |\Gamma|=n} R(\Gamma; q), \\ \bar{D}_n(q) &= \sum_{\Gamma \in \mathcal{F}^\gamma, |\Gamma|=n} R(\Gamma; q),\end{aligned}$$

and similarly for the self-energy and the vacuum polarization. The renormalized propagators $\bar{S}(q; e)$ and $\bar{D}(q; e)$ can then be described as power series over the square of the (finite) effective charge e ,

$$\bar{S}(q; e) = \sum_{n \geq 0} e^{2n} \bar{S}_n(q), \tag{7}$$

$$\bar{D}(q; e) = \sum_{n \geq 0} e^{2n} \bar{D}_n(q). \tag{8}$$

In the simplest case, e is the charge of the electron experimentally measured.

In a strike of genius, Freeman Dyson [13] noticed that the relation between the renormalized propagators $\bar{S}(q; e)$, $\bar{D}(q; e)$ and the bare propagators $S(q; e_0)$, $D(q; e_0)$ can be given by introducing two (scalar) renormalization factors $Z_2(e)$ and $Z_3(e)$ which withdraw the divergent part of $S(q; e_0)$ and $D(q; e_0)$ according to the equations

$$Z_2(e) \bar{S}(q, e) = S(q, e_0), \tag{9}$$

$$Z_3(e) \bar{D}(q, e) = D(q, e_0). \tag{10}$$

Equations (9) and (10) are known as Dyson's formulas. J.C. Ward then showed [30] that the renormalized charge e is related to the bare charge e_0 , which is an infinite value, by means of the photon renormalization factor,

$$Z_3(e)^{-1/2}e = e_0. \quad (11)$$

In modern textbooks, cf. [31, 27], the renormalization factors Z_2 and Z_3 are introduced from the ratio between renormalized and bare fields, from the counterterms of the Lagrangian, from the residue of the pole of the propagators, sometimes also from the field commutation relations. We will show that Dyson's formulas are very effective for an extension to nonscalar renormalization factors. In fact, in this paper we discuss how they determine some relations between the coefficients of the perturbative expansions, and finally how these relations determine a new noncommutative form of the Dyson's formulas (9,10) and of the charge renormalization (11).

Since the renormalization conditions are given via the self-energy and the vacuum polarization, which are expanded over 1PI Feynman diagrams, it is clear that the renormalization factors Z_2 and Z_3 are related to the counterterms $C(\Gamma)$ of 1PI diagrams [7].

In particular, to fix a notation, let us expand the renormalization factors over e^2 as

$$Z_2(e) = 1 + \sum_{n \geq 1} e^{2n} Z_{2,n}, \quad (12)$$

$$Z_3(e) = 1 - \sum_{n \geq 1} e^{2n} Z_{3,n}, \quad (13)$$

where $Z_{2,n}$ and $Z_{3,n}$ are the renormalization terms which appear at the n -th order of interaction. The minus sign in the definition of $Z_3(e)$ is chosen for later convenience (it is similar to the minus sign used in the Connes-Kreimer homomorphism [12]). Then, the coefficients $Z_{2,n}$ and $Z_{3,n}$ are related to the counterterms as follows, cf. [7],

$$Z_{2,n} = - \sum_{\Gamma \in \mathcal{F}_{1PI}^e, |\Gamma|=n} C_2(\Gamma; q),$$

$$Z_{3,n} = \sum_{\Gamma \in \mathcal{F}_{1PI}^\gamma, |\Gamma|=n} C_3(\Gamma; q).$$

One of the main features of Feynman diagrams is the fact that the amplitude of two graphs joined by a single line is computed as the product of the amplitudes of the disjoint graphs. In other words, the junction of adjacent Feynman diagrams defines an internal associative product on the sets $\mathcal{F}^e$ and $\mathcal{F}^\gamma$ compatible with the map U , with generators given by the subspaces $\mathcal{F}_{1PI}^e$ and $\mathcal{F}_{1PI}^\gamma$.

In fact, this assertion is not entirely correct. The propagators $S(q; e_0)$ and $D(q; e_0)$ of Eqs. (5,6) are described by graphs which start and end with a free propagator (a simple line). When we joint two such graphs, one of the two central lines of the new graph disappears, and this produces a new factor in the amplitudes, proportional to the inverse of the amplitude of the free propagator. For instance,

$$U(\text{---}\bigcirc\text{---}\bigcirc\text{---}) = U(\text{---}\bigcirc\text{---}) \frac{1}{U(\text{---})} U(\text{---}\bigcirc\text{---}).$$

To avoid this factor we choose a notation of Feynman diagrams which differs slightly from the usual one: we drop the first simple line from all diagrams. This modification of the diagrams corresponds to divide all propagators by the free propagator on the left, and to multiply the vacuum polarization and the self-energy by the free propagators on the right. In particular, the free propagators D^0 and S^0 become unit matrices and both correspond to no graph. We denote them by 1. The sets of Feynman diagrams

that we consider are then the following:

$$\begin{aligned}\mathcal{F}^e &= \{1, \text{cloud}, \text{cloud with loop}, \text{cloud with bubble}, \text{cloud with triangle}, \text{cloud with square}, \dots\} \\ \mathcal{F}^\gamma &= \{1, \text{fermion loop}, \text{fermion loop with cloud}, \text{fermion loop with bubble}, \text{fermion loop with triangle}, \text{fermion loop with square}, \dots\}\end{aligned}$$

With these notations, the map U becomes multiplicative, that is, $U(\Gamma_1)U(\Gamma_2) = U(\Gamma_1\Gamma_2)$. For instance,

$$U(\text{fermion loop with fermion loop}) = U(\text{fermion loop})U(\text{fermion loop})$$

and

$$U(\text{cloud with cloud}) = U(\text{cloud})U(\text{cloud}).$$

The 1PI Feynman diagrams are then the generators of the algebra of all diagrams with the “junction” product. Since no other relationship between the diagrams is forced by the amplitude map U , the algebra of Feynman diagrams can also be seen as the free algebra on the 1PI diagrams, that is,

$$\begin{aligned}(\mathcal{F}^\gamma, \text{junction}) &\cong (T(\mathcal{F}_{1PI}^\gamma), \otimes), \\ (\mathcal{F}^e, \text{junction}) &\cong (T(\mathcal{F}_{1PI}^e), \otimes).\end{aligned}$$

2 From Feynman graphs to planar binary trees

In [6], [7] we proposed another way to describe the coefficients $S_n(q)$ and $D_n(q)$ of Eqs. (5, 6). It is an intermediate step between the calculation of single Feynman diagrams and the value of the sums $S_n(q)$ and $D_n(q)$. The idea is to represent the sum of “similar” Feynman diagrams with a single symbol, namely a planar binary tree. The purpose is twofold: on one side we can make use of the rich algebraic structure of the set of trees to have new informations on the perturbative terms; on the other side we can still comfortably deal with the restricted sums of Feynman diagrams. For instance, when the fermions are heavy, it is convenient to limit the calculation to a small number of fermion loops. The main example is quenched QED, cf. [4], where the photon propagator is described by the Feynman diagrams without vacuum polarization insertions.

The choice of planar binary trees is motivated by the fact, showed in [5], that the functional Schwinger equations satisfied by the QED propagators have a very natural perturbative solution in terms of such trees. As we said, the expansion of the propagators over trees gives more detailed information than the sums $S_n(q)$ and $D_n(q)$. This situation is analogue to Jean-Louis Loday’s arithmetree [22], where each natural number n is written as the sum of all planar binary trees with n internal vertices, and operations are defined on trees, that are compatible with the addition and the product of integers.

Passing from Feynman graphs to trees has the disadvantage of losing the intuition of each ‘physical interaction’, but not that of ‘order of interaction’, since each leaf of the tree detects the presence of a loop in the corresponding graphs. The main advantage of the tree representation of interactions is that overlapping divergencies do not appear isolated, and their renormalization is encoded by the same procedure used for simple divergencies. Moreover, the Ward identities are built into the tree representation, so their consequences are automatically taken into account.

We remark, once more, that the present work is intended to deal only with the algebraic and combinatorial structure which encodes renormalization. In no way it simplifies the analytical computations of the amplitudes. In particular, in [5] and [7] we showed that the tree-expansion method has a peculiarity,

if compared to the graph-expansion one. Namely, the coefficients of the expansions satisfy an explicit recursive formula on the order of the trees, in both cases before and after the renormalization. However, as kindly remarked to the authors by D. Kreimer and D. Broadhurst, the difficulty arising in the analytical computation of each integral described by Feynman graphs is only postponed to the computation of the recursive terms described by trees. Moreover, it should be added that the renormalization in the tree representation does not produce a finite result for each tree, as explained in Ref.[7]. The tree representation is used only as a powerful method to generate recursive equations which, when summed over all trees of a given order, will give a finite result. These recursive equations are required to define the Hopf algebra of renormalization.

2.1 Correspondence between trees and graphs

Planar binary trees are connected planar graphs with trivalent internal vertices and no cycle. The external edges are called the leaves. We draw them with one preferred leaf on the bottom, that we call the root. Let Y denote the set of planar binary trees, simply called “trees”. Trees are naturally graded by the number of internal vertices. We denote by $|t|$ the number of internal vertices of the tree t , and by Y_n the set of trees t with $|t| = n$. Then, $Y = \bigcup_{n \geq 0} Y_n$, where $Y_0 = \{\text{leaf}\}$, $Y_1 = \{\text{trivalent vertex}\}$, $Y_2 = \{\text{two trivalent vertices}\}$, etc.

The correspondence between trees and Feynman’s diagrams can be chosen as two maps

$$\begin{aligned}\sigma^e : Y &\longrightarrow \mathcal{F}^e, \\ \sigma^\gamma : Y &\longrightarrow \mathcal{F}^\gamma,\end{aligned}$$

where $\mathcal{F}^e$ and $\mathcal{F}^\gamma$ now denote the complex vector spaces spanned by the sets of graphs, such that their image exhausts all the 2-legs Feynman diagrams of QED at a fixed order of interaction, that is

$$\begin{aligned}\sum_{t \in Y, |t|=n} \sigma^e(t) &= \sum_{\Gamma \in \mathcal{F}^e, |\Gamma|=n} \Gamma, \\ \sum_{t \in Y, |t|=n} \sigma^\gamma(t) &= \sum_{\Gamma \in \mathcal{F}^\gamma, |\Gamma|=n} \Gamma.\end{aligned}$$

Moreover, each graph appears once and only once as an addend in $\sigma^e(t)$ or $\sigma^\gamma(t)$ for some tree t . In this way, if we extend linearly the map U to formal sums of graphs, the Feynman rules can be read directly on trees, through the compositions

$$\varphi^i = U \circ \sigma^i, \quad \text{for } i = e, \gamma, \quad (14)$$

and the QED propagators can be developed as formal series over the trees, with amplitudes φ^i .


To define the maps σ^i , for $i = e, \gamma$, we follow some basic rules:

- The orientation “left to right” in a Feynman diagram corresponds to the orientation “root to leaves” in a tree.
- Electron propagators correspond to $/$ -branches; photon propagators correspond to $\backslash$ -branches.
- Electron loops correspond to $/$ -leaves; photon loops correspond to $\backslash$ -leaves.
- Subloops are loops of a subdiagram.

To make the next algorithms clear, we also fix a precise definition of the electron and photon loops we shall encounter:


- A photon loop in a 2-legs electron graph is a 2-legs photon graph such that if we remove it we still have a 2-legs electron graph. For instance, in the electron graph





if we remove , we remain with , which is not a 2-legs diagram. Therefore,

the only photon loop we can consider is the free propagator .


- An electron loop in a 2-legs photon graph is (the closure of) a 2-legs electron graph such that if we remove it we still have a 2-legs photon graph. For instance, in the photon graph



we can not remove , because it is the closure of a 3-legs diagram . Therefore, the only electron loop we can consider is the diagram



which is the closure of the 2-legs diagram



Starting from the Schwinger-Dyson equations for the propagators $S(q; e_0)$ and $D(q; e_0)$, translated into a recursive formula of perturbative amplitudes in [6], we are forced to define the maps σ^e and σ^γ through a recursive algorithm, which is better understood as the inverse of the following one.

Definition 2.1 Define the maps $\pi^e : \mathcal{F}^e \longrightarrow Y$ and $\pi^\gamma : \mathcal{F}^\gamma \longrightarrow Y$ recursively, starting from the free propagators $\pi^e(1), \pi^\gamma(1) = \mathbf{1}$. For any 2-legs Feynman diagram Γ , follow this algorithm. If Γ is an electron graph:

- Draw a $\backslash$ -leaf and read the graph from left to right.
- Graft a $\backslash$ -leaf at the bottom of the $\backslash$ -leaf for the first photon loop encountered.
- If the photon loop has subloops, draw a photon subtree following the algorithm below for photon graphs.
- Return to the first vertex of the photon loop and start the algorithm from the top of the rightmost $\backslash$ -leaf.
- Never read the second vertex of the photon loops. The algorithm ends when all the first vertices of the photon loops are met. At the end, draw the root.

If Γ is a photon graph:

- Draw a $\backslash$ -leaf and read the graph from left to right.
- Graft a $\backslash$ -leaf at the bottom of the $\backslash$ -leaf for the first electron loop encountered.

- If the electron loop has subloops, draw an electron subtree following the algorithm above for electron graphs.
- Return to the second vertex of the electron loop and start the algorithm from the top of the leftmost $\setminus$ -leaf.
- The algorithm ends when all the electron loops are met. At the end, draw the root.

As an example, step by step, let us compute $\pi^e(\text{cloud on line})$, knowing that $\pi^\gamma(\text{wavy}) = \pi^\gamma(1) = 1$:

$$\pi^e(\text{cloud on line}) : \quad / , \quad \vee , \quad \text{Y} .$$

Similarly, let us compute $\pi^\gamma(\text{circle on wavy})$, knowing that $\pi^e(\text{line}) = \pi^e(1) = 1$:

$$\pi^\gamma(\text{circle on wavy}) : \quad \setminus , \quad \vee , \quad \text{Y} .$$

More complicated examples:

$$\pi^e(\text{cloud on cloud on line}) : \quad / , \quad \vee , \quad \text{Y} \vee \text{Y} , \quad \text{Y} \vee \text{Y} \vee \text{Y} ;$$

$$\pi^e(\text{circle on cloud on line}) : \quad / , \quad \text{Y} \vee \text{Y} , \quad \text{Y} \vee \text{Y} \vee \text{Y} ;$$

$$\pi^\gamma(\text{circle on circle on wavy}) : \quad \setminus , \quad \text{Y} \vee \text{Y} , \quad \text{Y} \vee \text{Y} \vee \text{Y} ;$$

$$\pi^\gamma(\text{circle on circle on circle on wavy}) : \quad \setminus , \quad \vee , \quad \text{Y} \vee \text{Y} , \quad \text{Y} \vee \text{Y} \vee \text{Y} .$$

The maps π^e and π^γ are not injective, but they are surjective and “finite-to-one”. Therefore, they admit two linear sections σ^e and σ^γ (left inverse maps) which can be described, up to some scalar factors, as follows.

Let $\vee : Y_n \times Y_m \longrightarrow Y_{n+m+1}$ be the map which grafts two trees on a new root. Then, each tree t except the root tree 1 is the grafting of two trees with strictly smaller order,

$$\begin{aligned} t &= t^l \vee t^r = \vee(t^l, t^r), \\ |t| &= |t^l| + |t^r| + 1, \end{aligned}$$

the first one on the left and the second one on the right-hand side of the root. For instance,

$$\text{Y} \vee \text{Y} = \text{Y} \vee 1, \quad \text{Y} \vee \text{Y} = \text{Y} \vee \text{Y}.$$

Definition 2.2 Define the maps $\sigma^e : Y \longrightarrow \mathcal{F}^e$ and $\sigma^\gamma : Y \longrightarrow \mathcal{F}^\gamma$ recursively, starting from the free propagators $\sigma^e(1), \sigma^\gamma(1) = 1$. For any tree $t \neq 1$, proceed in two different ways to build $\sigma^e(t)$ and $\sigma^\gamma(t)$. For electron trees:

where the order of interaction is now given by the number $|t|$ of internal vertices of the tree t .

After the renormalization of each finite sum of integrals contained in $\varphi^e(t; q)$ and $\varphi^\gamma(t; q)$, we denote the finite parts by $\bar{\varphi}^e(t; q)$ and $\bar{\varphi}^\gamma(t; q)$. The coefficients $\bar{S}_n(q)$ and $\bar{D}_n(q)$ of the renormalized propagators are then given by

$$\bar{S}_n(q) = \sum_{|t|=n} \bar{\varphi}^e(t; q), \quad (17)$$

$$\bar{D}_n(q) = \sum_{|t|=n} \bar{\varphi}^\gamma(t; q). \quad (18)$$

The relationship between the coefficients of the tree-expansion before and after the massless renormalization was found in [7] by means of a Hopf algebra structure on the set of trees. We recall it at the end of section 3, where we introduce the necessary notations.

In the next two sections we define a multiplicative structure on the set of trees which is compatible with the (noncommutative) junction of adjacent 2-legs Feynman graphs of QED.

2.2 Algebra of photon trees and vacuum polarization

For photon trees, the Schwinger-Dyson equation for $D(q)$ provides the relation $\varphi^\gamma(t^l \vee t^r) = \varphi^\gamma(\iota \vee t^r) \varphi^\gamma(t^l)$ for the amplitudes (see Refs.[6, 7]). For example, applying the definition (2.1) of the map σ^γ , we find

$$\begin{aligned} \varphi^\gamma(\Upsilon; q) &= U(\text{diagram: a circle with a wavy line entering from the left and a wavy line exiting from the right}) \\ \varphi^\gamma(\check{\Upsilon}; q) &= U(\text{diagram: a circle with a wavy line entering from the left and a dashed line exiting from the right}) + \text{diagram: a circle with a dashed line entering from the left and a wavy line exiting from the right} + \text{diagram: a circle with a dashed line entering from the left and a dashed line exiting from the right}) \\ \varphi^\gamma(\check{\check{\Upsilon}}; q) &= U(\text{diagram: a circle with a wavy line entering from the left and a dashed line exiting from the right, with a dashed line loop on the left}) + \text{diagram: a circle with a dashed line entering from the left and a wavy line exiting from the right, with a dashed line loop on the right}) + \text{diagram: a circle with a dashed line entering from the left and a dashed line exiting from the right, with a dashed line loop on the top}) \end{aligned}$$

so we see that $\varphi^\gamma(\check{\check{\Upsilon}}; q) = \varphi^\gamma(\Upsilon; q) \varphi^\gamma(\check{\Upsilon}; q)$. We transform this relation between amplitudes $\varphi^\gamma(t; q)$ into a relation between trees.

Definition 2.3 Denote by $\mathcal{H}^\gamma = Y$ the graded vector space generated by Y (with $\mathcal{H}_n^\gamma = Y_n$), equipped with the graded product $\cdot : \mathcal{H}_n^\gamma \otimes \mathcal{H}_m^\gamma \longrightarrow \mathcal{H}_{n+m}^\gamma$ ¹, defined by the recurrence relation

$$\begin{aligned} s \cdot t &:= (s^l \cdot t) \vee s^r \quad \text{for } s = s^l \vee s^r, \\ \iota \cdot t &:= t. \end{aligned}$$

This product consists in grafting the tree t onto the leftmost leaf of tree s . Therefore, it is associative and has a unit given by the root tree ι . In particular, $(\iota \vee s) \cdot t = t \vee s$, which is the relation we wanted to implement. Examples of products are

$$\begin{aligned} \check{\Upsilon} \cdot \check{\Upsilon} &= \check{\check{\Upsilon}}, \\ \Upsilon \cdot \check{\Upsilon} &= \check{\check{\Upsilon}} = \check{\Upsilon} \cdot \Upsilon = \Upsilon \cdot \Upsilon, \\ \Upsilon \cdot \check{\check{\Upsilon}} &= \check{\check{\check{\Upsilon}}}, \quad \check{\Upsilon} \cdot \Upsilon = \check{\check{\Upsilon}}. \end{aligned}$$

The $\cdot$ product is defined in such a way that the following property automatically holds:

Lemma 2.4 *The map $\sigma^\gamma : \mathcal{H}^\gamma \longrightarrow \mathcal{F}^\gamma$ is an algebra morphism, that is, the $\cdot$ product between photon trees is transformed into the junction of adjacent Feynman graphs. Therefore, the amplitude map φ^γ (and $\bar{\varphi}^\gamma$) is also an algebra morphism on trees.*

¹ The product taken in $\mathcal{H}^\gamma$ is the opposite of the operation over defined by Loday and Ronco in [26].

Denote by $V : Y_n \longrightarrow Y_{n+1}$ the left grafting by the root,

$$V(t) = \mathfrak{r} \vee t. \quad (19)$$

Then, any tree t can be decomposed as

$$t = t^l \vee t^r = V(t^r) t^l,$$

where the left tree t^l can be further decomposed until we reach a root tree $\mathfrak{r}$ on the left. Therefore, the elements $V(t)$, for all $t \in Y$, form a system of generators for the algebra $\mathcal{H}^\gamma$.

Proposition 2.5 *Under the map σ^γ , the generators $V(t)$ of the algebra $\mathcal{H}^\gamma$ correspond to sums of 1PI 2-legs Feynman diagrams for the photon propagator.*

Therefore, the bare and renormalized vacuum polarizations are expanded only on the generator trees as

$$\Pi(q; e_0) = \sum_t e_0^{2|t|+2} \varphi^\gamma(V(t); q), \quad (20)$$

$$\bar{\Pi}(q; e) = \sum_t e^{2|t|+2} \bar{\varphi}^\gamma(V(t); q). \quad (21)$$

This fact is evident from the algorithm defining σ^γ in (2.2).

Lemma 2.6 *Any formal series in $\mathcal{H}^\gamma[[x]]$ starting with $\mathfrak{r}$ can be inverted, w.r.t. the product given in (2.3). Moreover the inverse series is developped only on generator trees,*

$$\left(\mathfrak{r} + \sum_{|t|>0} x^{|t|} t \right)^{-1} = \mathfrak{r} - \sum_t x^{|t|+1} V(t). \quad (22)$$

Proof. Since φ^γ is an algebra homomorphism, formula (22) could be deduced from the equation (1). But we give here a direct proof, based on the typical property of trees, cf. [25, 15],

$$\sum_{|t|=n} t = \sum_{k=0}^{n-1} \sum_{|t^l|=k} \sum_{|t^r|=n-k-1} t^l \vee t^r, \quad (23)$$

valid for $n > 0$. We obtain

$$\begin{aligned} \mathfrak{r} + \sum_{|t|>0} x^{|t|} t &= \mathfrak{r} + \sum_{t^l} \sum_{t^r} x^{|t^l|+|t^r|+1} t^l \vee t^r = \mathfrak{r} + \sum_{t^l} \sum_{t^r} x^{|t^l|+|t^r|+1} V(t^r) t^l \\ &= \mathfrak{r} + \left(\sum_t x^{|t|+1} V(t) \right) \left(\mathfrak{r} + \sum_{|t|>0} x^{|t|} t \right). \end{aligned}$$

Hence,

$$\left(\mathfrak{r} - \sum_t x^{|t|+1} V(t) \right) \left(\mathfrak{r} + \sum_{|t|>0} x^{|t|} t \right) = \mathfrak{r}.$$

□

² In fact, no other relation than associativity holds between any product of generators, hence $\mathcal{H}^\gamma$ is isomorphic to the free associative unital algebra $\mathbb{C}\langle Y \rangle$ generated by the set of trees, where the unit $1 \in \mathbb{C}$ corresponds to the root tree $\mathfrak{r}$.

2.3 Algebra of electron trees and self-energy

The algebra of trees for electrons is a little more subtle, because no particular subset of trees corresponds to 1PI graphs under the map σ^e , and no internal product analogue to $\circ$ is forced from the Schwinger-Dyson equations. For instance, if $\varphi^e(\Upsilon)$ is a contribution to the electron propagator, $\varphi^e(\Upsilon)\varphi^e(\Upsilon)$ cannot be represented as $\varphi^e(t)$ for a tree t . Therefore, the product is not given by an internal low, and we have to consider the free (tensor) product.

If we want the map σ^e to be an algebra morphism, however, the set of generators for the electron algebra must correspond to 1PI electron graphs. Therefore, we start by describing the 1PI 2-legs Feynman diagrams for the electron propagator, using the pruning operator P introduced in [6]. We recall its definition and main properties.

Definition 2.7 Define an operator $\Delta^P : Y \longrightarrow Y \otimes Y$ recursively as

$$\begin{aligned}\Delta^P \mathbf{1} &= \mathbf{1} \otimes \mathbf{1}, \\ \Delta^P(s \vee t) &= \mathbf{1} \otimes (s \vee t) + (\vee \otimes \text{Id})(s \otimes \Delta^P t),\end{aligned}\tag{24}$$

where $\vee(s, t) = s \vee t$. For example:

$$\begin{aligned}\Delta^P \Upsilon &= \mathbf{1} \otimes \Upsilon + \Upsilon \otimes \mathbf{1}, \\ \Delta^P \check{\Upsilon} &= \mathbf{1} \otimes \check{\Upsilon} + \Upsilon \otimes \Upsilon + \check{\Upsilon} \otimes \mathbf{1}, \\ \Delta^P \check{\check{\Upsilon}} &= \mathbf{1} \otimes \check{\check{\Upsilon}} + \check{\Upsilon} \otimes \mathbf{1}.\end{aligned}$$

The pruning operator of [6] is then $P(t) = \Delta^P(t) - t \otimes \mathbf{1} - \mathbf{1} \otimes t$. In ref. [7], it was shown that Δ^P is coassociative. Moreover, the coproduct Δ^P preserves the order of the trees, in fact $\Delta^P(Y_n) \subset \bigoplus_{p+q=n} Y_p \otimes Y_q$. Therefore, if we extend it multiplicatively on tensor product of trees, it is easy to see that the operator Δ^P defines a graded Hopf algebra over $\mathbb{C}\langle Y \rangle / (\mathbf{1} - 1)$, with respect to the grading given by the total order of products of trees, that is

$$|t_1 \otimes \cdots \otimes t_k| = |t_1| + \cdots + |t_k|.$$

The counit is the algebra morphism defined on generators as $\epsilon(t) = 0$ for all $t \neq \mathbf{1}$ and $\epsilon(\mathbf{1}) = 1$, and the pruning antipode S_P is the (graded) algebra anti-morphism automatically defined on generators by the recursive formula $S_P(\mathbf{1}) = \mathbf{1}$ and

$$S_P(t) = -t - \sum_{P(t)} S_P(t_{(1)})t_{(2)} = -t - \sum_{P(t)} t_{(1)}S_P(t_{(2)}),$$

where we intentionally omit the tensor product symbol between trees. We give a few examples:

$$\begin{aligned}S_P(\Upsilon) &= -\Upsilon, \\ S_P(\check{\Upsilon}) &= -\check{\Upsilon} + \Upsilon^2, \\ S_P(\check{\check{\Upsilon}}) &= -\check{\check{\Upsilon}}, \\ S_P(\check{\check{\check{\Upsilon}}}) &= -\check{\check{\check{\Upsilon}}} + \check{\Upsilon}\Upsilon + \Upsilon\check{\Upsilon} - \Upsilon^3, \\ S_P(\check{\check{\check{\check{\Upsilon}}}}) &= -\check{\check{\check{\check{\Upsilon}}}} + \Upsilon\check{\check{\Upsilon}}.\end{aligned}$$

Notice that the coproduct Δ^P is neither commutative nor cocommutative, and $S_P^2 \neq \text{Id}$.

Remark 2.8 An alternative way to define Δ^P was pointed out to us by J.-L. Loday. In Ref.[26], the operation *under* was defined recursively by $\mathbf{1} \setminus t = t$ and $s \setminus t = s^l \vee (s^r \setminus t)$ for $s = s^l \vee s^r$. Then the coproduct Δ^P is the dual operation to the product *under*, in the sense that

$$\Delta^P t = t \otimes \mathbf{1} + \mathbf{1} \otimes t + \sum_{s_1 \setminus s_2 = t} s_1 \otimes s_2,$$

with the convention that the sum is over all $s_1 \neq \mathbf{1}$, $s_2 \neq \mathbf{1}$ such that $s_1 \setminus s_2 = t$. If t cannot be written as $s_1 \setminus s_2$ for $s_1 \neq \mathbf{1}$ and $s_2 \neq \mathbf{1}$, then $\Delta^P t = t \otimes \mathbf{1} + \mathbf{1} \otimes t$.

Lemma 2.9 *Any formal series in $\mathbb{C}\langle Y \rangle / (\imath - 1)[[x]]$ starting with $\imath$ can be inverted, w.r.t. the tensor product. Moreover the inverse series is developed only on trees of the form $S_P(t)$, with $t \in Y$, that is,*

$$\left(\imath + \sum_{|t|>0} x^{|t|} t\right)^{-1} = \imath + \sum_{|t|>0} x^{|t|} S_P(t). \quad (25)$$

Proof. The main property of Δ^P was noticed in Ref.[6]:

$$\begin{aligned} \left(\imath + \sum_{|t|>0} x^{|t|} t\right) \otimes \left(\imath + \sum_{|t|>0} x^{|t|} t\right) &= \imath \otimes \imath + \sum_{|t|>0} x^{|t|} (t \otimes \imath + \imath \otimes t + \sum_{P(t)} t_{(1)} \otimes t_{(2)}) \\ &= \imath \otimes \imath + \sum_{|t|>0} x^{|t|} \Delta^P t. \end{aligned} \quad (26)$$

From Eq. (26) and the definition of the antipode we obtain

$$\left(\imath + \sum_{|t|>0} x^{|t|} S_P(t)\right) \left(\imath + \sum_{|t|>0} x^{|t|} t\right) = \imath + \sum_{|t|>0} x^{|t|} (S_P(t) + t + \sum_{s_1 \setminus s_2 = t} S_P(s_1) s_2) = \imath.$$

□

By definition (2) of the self-energy, and knowing that it should be expanded on 1PI Feynman graphs, we obtain:

Proposition 2.10 *The bare and renormalized self-energy are expanded as*

$$\begin{aligned} \Sigma(q; e_0) &= - \sum_{|t|>0} e_0^{2|t|} \varphi^e(S_P(t); q), \\ \bar{\Sigma}(q; e) &= - \sum_{|t|>0} e^{2|t|} \bar{\varphi}^e(S_P(t); q). \end{aligned}$$

Therefore, if we extend multiplicatively the map $\sigma^e : Y \longrightarrow \mathcal{F}^e$ to tensor powers of Y , the elements $S_P(t) \in \mathbb{C}\langle Y \rangle / (\imath - 1)$ correspond to sums of 1PI 2-legs Feynman diagrams for the electron propagator.

Now we know which must be the generators for the electron algebra.

Definition 2.11 We define $\mathcal{H}^e = \mathbb{C}\langle S_P(Y) \rangle / (\imath - 1)$ as the free associative algebra on the set of trees transformed by S_P , where we identify the formal unit 1 with the root tree $\imath$. For notational convenience, we omit the tensor product symbols in the algebra $\mathcal{H}^e$.

The vector space of planar binary trees Y can be seen as a subspace of $\mathcal{H}^e$, because by definition of the antipode S_P each tree $t \in Y$ can be seen as a (noncommutative) polynomial

$$t = -S_P(t) - \sum_{P(t)} S_P(t_{(1)}) t_{(2)},$$

where $t_{(2)}$ can be further developed as a polynomial in variables $S_P(t')$. Therefore the two algebras $\mathbb{C}\langle Y \rangle / (\imath - 1)$ and $\mathbb{C}\langle S_P(Y) \rangle / (\imath - 1) = \mathcal{H}^e$ are isomorphic and the map $\sigma^e : Y \longrightarrow \mathcal{F}^e$ can be extended as an algebra morphism to the space $\mathcal{H}^e$.

Remark 2.12 Beside the natural grading coming from the tensor powers, on a product of electron trees we can define a total order as the sum of the orders of the trees,

$$|t_1 \cdots t_k| = |t_1| + \cdots + |t_k|.$$

Then the algebra $\mathcal{H}^e$ of electron trees can be decomposed into homogeneous components

$$\mathcal{H}_n^e = \bigoplus_{n_1 + \cdots + n_k = n} S_P(Y_{n_1}) \otimes \cdots \otimes S_P(Y_{n_k}),$$

and since the pruning antipode is a homogeneous operator, these components can be written as

$$\mathcal{H}_n^e = \bigoplus_{n_1 + \dots + n_k = n} Y_{n_1} \otimes \dots \otimes Y_{n_k}.$$

To end this section, we give an example in terms of Feynman diagrams. Let us draw the perturbative expansion of $S(q; e_0)$ and $\Sigma(q; e_0)$ up to order 2. The propagator is

$$S(q; e_0) = 1 + e_0^2 \varphi^e(\Upsilon; q) + e_0^4 \varphi^e(\check{\Upsilon}; q) + e_0^4 \varphi^e(\check{\check{\Upsilon}}; q) + \mathcal{O}(e_0^6),$$

where

$$\begin{aligned} \varphi^e(\Upsilon) &= U\left(\text{diagram: a horizontal line with a wavy loop on top} \right) \\ \varphi^e(\check{\Upsilon}) &= U\left(\text{diagram: a horizontal line with a wavy loop on top, wavy line on bottom} \right) + U\left(\text{diagram: a horizontal line with a wavy loop on top, wavy line on bottom, wavy line on top} \right) + U\left(\text{diagram: a horizontal line with two wavy loops on top} \right), \\ \varphi^e(\check{\check{\Upsilon}}) &= U\left(\text{diagram: a horizontal line with a circle loop on top} \right). \end{aligned}$$

Therefore, the self-energy is now written as

$$\Sigma(q; e_0) = -e_0^2 \varphi^e(S_P(\Upsilon); q) - e_0^4 \varphi^e(S_P(\check{\Upsilon}); q) - e_0^4 \varphi^e(S_P(\check{\check{\Upsilon}}); q) - \mathcal{O}(e_0^6),$$

where

$$\begin{aligned} \varphi^e(S_P(\Upsilon)) &= -\varphi^e(\Upsilon) = -U\left(\text{diagram: a horizontal line with a wavy loop on top} \right) \\ \varphi^e(S_P(\check{\Upsilon})) &= -\varphi^e(\check{\Upsilon}) + \varphi^e(\Upsilon)\varphi^e(\Upsilon) = -U\left(\text{diagram: a horizontal line with a wavy loop on top, wavy line on bottom} \right) - U\left(\text{diagram: a horizontal line with a wavy loop on top, wavy line on bottom, wavy line on top} \right), \\ \varphi^e(S_P(\check{\check{\Upsilon}})) &= -\varphi^e(\check{\check{\Upsilon}}) = -U\left(\text{diagram: a horizontal line with a circle loop on top} \right). \end{aligned}$$

2.4 Gradings and loops

By construction (2.1), in any planar binary tree each leaf (except one!) corresponds to a loop in the associated Feynman diagrams. Photon and electron loops are distinguished through the orientation of the corresponding leaf: leaves growing to the right (i.e. /-leaves) correspond to electron loops, leaves growing to the left (i.e. \-leaves) correspond to photon loops. The remaining leaf is just a prolongation of the root, and we can identify it with the leftmost one for photon trees, and with the rightmost one for electron trees.

For any tree t , we denote by $L^e(t)$ and $L^\gamma(t)$ the number of electron and photon loops respectively. Since in planar binary trees each leaf (except one) corresponds to an internal vertex, we have exactly

$$L^e(t) + L^\gamma(t) = |t|.$$

For instance, in quenched QED, the photon trees t have $L^\gamma(t) = |t| - 1$, $L^e(t) = 1$, and the electron trees s have $L^\gamma(s) = |s|$, $L^e(s) = 0$.

To identify the space of photon and electron trees with a fixed number of photon or electron loops, we recall the double decomposition of the set of trees first introduced in [15].

For any $p, q \geq 0$, let $Y_{p,q}$ be the set of trees with p leaves oriented like $\backslash$ (except the leftmost one) and q leaves oriented like $/$ (except the rightmost one). Hence, the order of a (p, q) -tree t is $|t| = p + q + 1$, and the set of trees with order $n > 0$ can be decomposed as

$$Y_n = \bigoplus_{p+q+1=n} Y_{p,q}, \quad (27)$$

while Y_0 can not be decomposed.

Definition 2.13 The set of photon trees with no photon loops and no electron loops is, of course, Y_0 . Since a photon tree can not have any photon loop if there is no electron loop which supports it, we can single out the set of photon trees t with $L^\gamma(t) = p$ and $L^e(t) = q$ for any $p \geq 0$ and any $q > 0$ as the set $Y_{p,q-1}$. Therefore we define a loops bigrading on the vector space $\mathcal{H}^\gamma$ as

$$\begin{aligned} \mathcal{H}_{0,0}^\gamma &= Y_0, \\ \mathcal{H}_{p,0}^\gamma &= \emptyset, \quad \text{for any } p > 0, \\ \mathcal{H}_{p,q}^\gamma &= Y_{p,q-1}, \quad \text{for } p \geq 0 \text{ and } q > 0. \end{aligned}$$

It is easy to see that this bigrading yields a complete decomposition $\mathcal{H}_n^\gamma = \bigoplus_{p+q=n} \mathcal{H}_{p,q}^\gamma$, which is compatible with the product, that is

$$\mathcal{H}_{p_1,q_1}^\gamma \mathcal{H}_{p_2,q_2}^\gamma \subset \mathcal{H}_{p_1+p_2,q_1+q_2}^\gamma.$$

Similarly, the set of electron trees with no loops at all is Y_0 . Since on electron trees any electron loop needs at least one photon loop as support, this time we can identify the set of electron trees s with $L^\gamma(s) = p$ and $L^e(s) = q$ for any $p > 0$ and $q \geq 0$ as the set $Y_{p-1,q}$. In order to have a consistent extension of the loops bigrading with the free product of electron trees we define it as

$$\begin{aligned} \mathcal{H}_{0,0}^e &= Y_0, \\ \mathcal{H}_{0,q}^e &= \emptyset, \quad \text{for any } q > 0, \\ \mathcal{H}_{p,q}^e &= \bigoplus_{k=1}^p \bigoplus_{\substack{p_1+\dots+p_k=p \\ q_1+\dots+q_k=q}} \bigoplus_{q_1+\dots+q_k=q} Y_{p_1-1,q_1} \otimes \dots \otimes Y_{p_k-1,q_k}, \quad \text{for } p > 0 \text{ and } q \geq 0. \end{aligned}$$

Since the pruning antipode preserve the number of loops, to have an algebra decomposition the above vector space can be written as

$$\mathcal{H}_{p,q}^e = \bigoplus_{k=1}^p \bigoplus_{\substack{p_1+\dots+p_k=p \\ q_1+\dots+q_k=q}} S_P(Y_{p_1-1,q_1}) \otimes \dots \otimes S_P(Y_{p_k-1,q_k}).$$

Then at each order we have a complete decomposition $\mathcal{H}_n^e = \bigoplus_{p+q=n} \mathcal{H}_{p,q}^e$.

Remark 2.14 In [23], J.-L. Loday introduced a new type of associative algebra, called dialgebras, and their Koszul duals, simply called dual dialgebras. He then showed that the set of planar binary trees can be interpreted as the free dual dialgebra generated by one element. Therefore, in virtue of the operadic homology introduced by V. Ginzburg and M. Kapranov in [16], the set Y appears naturally in the dialgebra homology HY_* (moreover the order n of trees determines the degree of the homology, i.e. HY_n). Loday showed that the boundary operator of this homology is very elegantly expressed in terms of trees: it acts by deleting the leaves (see [24]).

More precisely, Loday showed that the dialgebra boundary operator is in fact simplicial, and each i -th face acts by deleting the i -th leaf of the trees. In [15], the simplicial complex on the set Y was recognized as the total complex of a bisimplicial structure, with horizontal and vertical face maps acting by deleting $/$ and $\backslash$ -leaves, respectively. The resulting bicomplex was shown to be acyclic.

Now, translating these results in terms of QED propagators, we obtain an acyclic bicomplex on the set of 2-legs Feynman diagrams, whose horizontal and vertical boundary operators act by deleting respectively electron and photon loops. How this can be interpreted in quantum field theory is an open question.

Remark 2.15 It is known that the number of trees with order n is the Catalan number $c_n = (2n)!/n!(n+1)!$. The number of trees with order n and a fixed quantity of photon loops p and of electron loops q has been computed in [15], as

$$\begin{aligned} c_{p,q}^\gamma &= \frac{(p+q+1)!(p+q+2)!}{p!(p+1)!(q-1)!q!} && \text{for photon trees,} \\ c_{p,q}^e &= \frac{(p+q+1)!(p+q+2)!}{(p-1)!p!q!(q+1)!} && \text{for electron trees.} \end{aligned}$$

There is an obvious symmetry $c_{p,q}^\gamma = c_{q,p}^e$.

3 Hopf algebras on planar binary trees

It is nowadays established that the relationship between the amplitudes of Feynman diagrams before and after the renormalization has a combinatorial description in terms of a Hopf algebra structure defined on the set of diagrams, cf. [19].

In [7], we showed that on the sets of trees there exists a structure of a Hopf algebra which encodes the renormalization of QED propagators. Looking at trees as sums of Feynman diagrams, the coproduct on trees is much simpler than the sum of the coproducts on each Feynman diagram. We pointed out that the collections of amplitudes described by each tree do not yield a finite value, but the sum of amplitudes at each order of interaction, renormalized through the tree coproduct, indeed give the usual value obtained by summing up the renormalized amplitudes of all Feynman diagrams, with a procedure which automatically encodes the Ward identities.

In this section we give the precise definition of the Hopf algebras defined on photon and electron trees, as quoted in [7]. Using these structures, in section 4 we will find the explicit formulas for the renormalization of QED propagators at each order of interaction.

3.1 Hopf algebra for the photon trees

Definition 3.1 Define a coproduct $\Delta^\gamma : \mathcal{H}^\gamma \longrightarrow \mathcal{H}^\gamma \otimes \mathcal{H}^\gamma$ by the recurrence relations

$$\begin{aligned} \Delta^\gamma \mathbf{1} &= \mathbf{1} \otimes \mathbf{1}, \\ \Delta^\gamma V(t) &= V(t) \otimes \mathbf{1} + \sum_{\Delta^\gamma t} t_{(1)} \otimes V(t_{(2)}) - \sum_{\Delta^\gamma t^l} (t_{(1)}^l \vee t^r) \otimes V(t_{(2)}^l) \\ &= V(t) \otimes \mathbf{1} + (\text{Id} \otimes V) \Delta^\gamma t - (\text{Id} \otimes V) [(V(t^r) \otimes \mathbf{1}) \Delta^\gamma t^l], \\ \Delta^\gamma (t \vee s) &= \Delta^\gamma V(s) \Delta^\gamma t, \end{aligned} \tag{28}$$

where we adopt Sweedler's convention $\Delta^\gamma t = \sum_{\Delta^\gamma t} t_{(1)} \otimes t_{(2)}$ and where $t = t^l \vee t^r$.

Let $\epsilon : \mathcal{H}^\gamma \longrightarrow \mathbb{C}$ be the map which sends all the trees to 0 except the root $\mathbf{1}$ which is sent to 1.

A few examples might be useful:

$$\begin{aligned} \Delta^\gamma \Upsilon &= \Upsilon \otimes \mathbf{1} + \mathbf{1} \otimes \Upsilon, \\ \Delta^\gamma \check{\Upsilon} &= \check{\Upsilon} \otimes \mathbf{1} + \mathbf{1} \otimes \check{\Upsilon} + 2\Upsilon \otimes \Upsilon, \\ \Delta^\gamma \check{\check{\Upsilon}} &= \check{\check{\Upsilon}} \otimes \mathbf{1} + \mathbf{1} \otimes \check{\check{\Upsilon}}, \\ \Delta^\gamma \check{\check{\check{\Upsilon}}} &= \check{\check{\check{\Upsilon}}} \otimes \mathbf{1} + \mathbf{1} \otimes \check{\check{\check{\Upsilon}}} + 3\Upsilon \otimes \check{\check{\Upsilon}} + 3\check{\check{\Upsilon}} \otimes \Upsilon, \\ \Delta^\gamma \check{\check{\check{\check{\Upsilon}}}} &= \check{\check{\check{\check{\Upsilon}}}} \otimes \mathbf{1} + \mathbf{1} \otimes \check{\check{\check{\check{\Upsilon}}}} + \Upsilon \otimes \check{\check{\check{\Upsilon}}} + \check{\check{\check{\Upsilon}}} \otimes \Upsilon, \\ \Delta^\gamma \check{\check{\check{\check{\check{\Upsilon}}}}} &= \check{\check{\check{\check{\check{\Upsilon}}}}} \otimes \mathbf{1} + \mathbf{1} \otimes \check{\check{\check{\check{\check{\Upsilon}}}}} + \Upsilon \otimes \check{\check{\check{\check{\Upsilon}}}} + \check{\check{\check{\check{\Upsilon}}}} \otimes \Upsilon, \\ \Delta^\gamma \check{\check{\check{\check{\check{\check{\Upsilon}}}}}} &= \check{\check{\check{\check{\check{\check{\Upsilon}}}}}} \otimes \mathbf{1} + \mathbf{1} \otimes \check{\check{\check{\check{\check{\check{\Upsilon}}}}}}. \end{aligned}$$

Proposition 3.2 *The space $\mathcal{H}^\gamma$ is a non-commutative nor cocommutative bigraded Hopf algebra.*

Proof. We first observe that the coproduct preserves the bigrading of $\mathcal{H}^\gamma$, that is,

$$\Delta^\gamma(\mathcal{H}_{p,q}^\gamma) \subset \bigoplus_{p_1+p_2=p, q_1+q_2=q} \mathcal{H}_{p_1,q_1}^\gamma \otimes \mathcal{H}_{p_2,q_2}^\gamma,$$

and therefore it also preserves the total order of trees defined in (2.3).

By recursion arguments, it is easy to see that the only terms of total order $(n, 0)$ and $(0, n)$, in the image of Δ^γ , are given by the primitive part $t \otimes \mathbf{1}$ and $\mathbf{1} \otimes t$ of any tree t . Then, the map ϵ is a counit for Δ^γ . Moreover (see ref. [29] for details), the antipode $S^\gamma : \mathcal{H}^\gamma \longrightarrow \mathcal{H}^\gamma$ is automatically defined by the recursive formula

$$S^\gamma(t) = -t - (S^\gamma \bar{\star} \text{Id})(t) \quad (29)$$

which involves the convolution $\bar{\star}$ of the reduced coproduct $\bar{\Delta}^\gamma(t) = \Delta^\gamma(t) - t \otimes \mathbf{1} - \mathbf{1} \otimes t$.

To prove that Δ^γ is coassociative, it is useful to introduce another operator $F : \mathcal{H}^\gamma \longrightarrow \mathcal{H}^\gamma \otimes \mathcal{H}^\gamma$, such that

$$\Delta^\gamma V(t) = V(t) \otimes \mathbf{1} + F(V(t)). \quad (30)$$

We achieve this by defining F again with some recurrence conditions

$$\begin{aligned} F(\mathbf{1}) &= \mathbf{1} \otimes \mathbf{1}, \\ F(V(t)) &= (\text{Id} \otimes V)F(t) = \sum_{F(t)} t_{(1)} \otimes V(t_{(2)}), \end{aligned}$$

and supposing that it is a multiplicative map, in the sense that

$$F(t \vee s) = F(V(s)) \Delta^\gamma t = \sum_{\Delta^\gamma(t), F(s)} (s_{(1)} t_{(1)}) \otimes t_{(2)} \vee s_{(2)}. \quad (31)$$

In fact, if we want Δ^γ to be given by Eq. (30), the operator F must be defined in such a way that

$$\begin{aligned} F(V(t)) &= \sum_{\Delta^\gamma t} t_{(1)} \otimes V(t_{(2)}) - \sum_{\Delta^\gamma t^l} (t_{(1)}^l \vee t^r) \otimes V(t_{(2)}^l) \\ &= (\text{Id} \otimes V) [\Delta^\gamma t - (V(t^r) \otimes \mathbf{1}) \Delta^\gamma t^l]. \end{aligned}$$

Therefore, if on a generator $V(t)$ we assume that $F(V(t)) = (\text{Id} \otimes V)F(t)$, we have

$$\begin{aligned} F(t) &= \Delta^\gamma t - (V(t^r) \otimes \mathbf{1}) \Delta^\gamma t^l \\ &= [\Delta^\gamma V(t^r) - (V(t^r)) \otimes \mathbf{1}] \Delta^\gamma t^l \\ &= F(V(t^r)) \Delta^\gamma t^l. \end{aligned}$$

First we prove by induction that the operator F defines a left Δ^γ -coaction of $\mathcal{H}^\gamma$ on itself, that is, it satisfies $(\text{Id} \otimes F)F = (\Delta^\gamma \otimes \text{Id})F$. Since F is multiplicative, we only need to show it on the generators $V(t)$. It is true on $t = \mathbf{1}$. Suppose that it is true for all the trees up to order n , and let $V(t)$ has order $n+1$. On one side we have

$$(\text{Id} \otimes F)F(V(t)) = (\text{Id} \otimes F \circ V)F(t) = (\text{Id} \otimes \text{Id} \otimes V)(\text{Id} \otimes F)F(t),$$

on the other side we have

$$(\Delta^\gamma \otimes \text{Id})F(V(t)) = (\Delta^\gamma \otimes V)F(t) = (\text{Id} \otimes \text{Id} \otimes V)(\Delta^\gamma \otimes \text{Id})F(t).$$

So the equality holds by inductive hypothesis.

Now we prove by induction that the operator Δ^γ is coassociative, using the fact that F is a coaction. Since Δ^γ is multiplicative, we only need to prove it on the generators $V(t)$. It is true on $t = \mathbf{1}$. Suppose

that Δ^γ is coassociative on all the trees with order up to n , and let $V(t)$ be a generator with order $n+1$. Then by the expression (30) of Δ^γ we have on one side

$$\begin{aligned} (\Delta^\gamma \otimes \text{Id})\Delta^\gamma V(t) &= \Delta^\gamma V(t) \otimes \mathbf{1} + (\Delta^\gamma \otimes \text{Id})F(V(t)) \\ &= V(t) \otimes \mathbf{1} \otimes \mathbf{1} + F(V(t)) \otimes \mathbf{1} + (\Delta^\gamma \otimes \text{Id})F(V(t)), \end{aligned}$$

and on the other side

$$\begin{aligned} (\text{Id} \otimes \Delta^\gamma)\Delta^\gamma V(t) &= V(t) \otimes \Delta^\gamma(\mathbf{1}) + (\text{Id} \otimes \Delta^\gamma)F(V(t)) \\ &= V(t) \otimes \mathbf{1} \otimes \mathbf{1} + (\text{Id} \otimes \Delta^\gamma \circ V)F(t) \\ &= V(t) \otimes \mathbf{1} \otimes \mathbf{1} + (\text{Id} \otimes V \otimes \text{Id})F(t) + (\text{Id} \otimes F \circ V)F(t) \\ &= V(t) \otimes \mathbf{1} \otimes \mathbf{1} + (\text{Id} \otimes V)F(t) \otimes \mathbf{1} + (\text{Id} \otimes F)(\text{Id} \otimes V)F(t) \\ &= V(t) \otimes \mathbf{1} \otimes \mathbf{1} + F(V(t)) \otimes \mathbf{1} + (\text{Id} \otimes F)F(V(t)). \end{aligned}$$

Then, the two sides are equal if and only if $(\Delta^\gamma \otimes \text{Id})F(t) = (\text{Id} \otimes F)F(t)$, which holds by inductive hypothesis. $\square$

3.2 Hopf structure for the electron trees

On the electron algebra $\mathcal{H}^e$, the coproduct operator analogue to the one on $\mathcal{H}^\gamma$ is not an internal operation. To describe it, we have to extend it to a bigger space which describes at the same time the electron and the photon propagators of QED.

Consider the tensor algebra $\mathcal{H}^{\text{qed}} = \mathcal{H}^\gamma \otimes \mathcal{H}^e$ with product defined componentwise as the product given in (28) on the photon trees and the tensor product on the electron trees. The unit is $\mathbf{1} \otimes \mathbf{1}$, and the two identifications $\mathcal{H}^\gamma = \mathcal{H}^\gamma \otimes \mathbf{1}$ and $\mathcal{H}^e = \mathbf{1} \otimes \mathcal{H}^e$ allow us to see $\mathcal{H}^\gamma$ and $\mathcal{H}^e$ as subalgebras of $\mathcal{H}^{\text{qed}}$. For the simplicity of notation, we denote by couples the elements of the tensor product between $\mathcal{H}^\gamma$ and $\mathcal{H}^e$. Hence we write $1 = (\mathbf{1}, \mathbf{1})$ for the unit $\mathbf{1} \otimes \mathbf{1} \in \mathcal{H}^{\text{qed}}$ and $(t, s_1 \dots s_n)$ for $t \otimes (s_1 \dots s_n) \in \mathcal{H}^\gamma \otimes \mathcal{H}^e$.

On the algebra $\mathcal{H}^{\text{qed}}$ we chose a grading given by the sum of the orders of all the trees appearing in a monomial. Hence, $\mathcal{H}^{\text{qed}}$ can be written as a direct sum $\mathcal{H}^{\text{qed}} = \bigoplus_{n \geq 0} \mathcal{H}_n^{\text{qed}}$ of homogeneous components $\mathcal{H}_n^{\text{qed}} = \bigoplus_{p+q=n} \mathcal{H}_p^\gamma \otimes \mathcal{H}_q^e$, where $\mathcal{H}_p^\gamma$ was defined in (2.3) and $\mathcal{H}_q^e$ in (2.12).

Definition 3.3 We define a coproduct $\Delta^{\text{qed}} : \mathcal{H}^{\text{qed}} \longrightarrow \mathcal{H}^{\text{qed}} \otimes \mathcal{H}^{\text{qed}}$ by its action on the photon and electron trees, and then we extend it on products as a morphism of algebras, that is

$$\begin{aligned} \Delta^{\text{qed}} 1 &= 1 \otimes 1, \\ \Delta^{\text{qed}}(t, s_1 \dots s_n) &= \Delta^\gamma(t) \Delta^e(s_1) \dots \Delta^e(s_n). \end{aligned}$$

On the photon trees, Δ^{qed} agrees with the coproduct Δ^γ defined above. On the electron trees, we define the coproduct $\Delta^e : \mathcal{H}^e \longrightarrow \mathcal{H}^{\text{qed}} \otimes \mathcal{H}^e$ (notice the photon component on the lefthand side!) by the recurrence relations

$$\begin{aligned} \Delta^e \mathbf{1} &= 1 \otimes \mathbf{1}, \\ \Delta^e(t \vee s) &= (\mathbf{1}, t \vee s) \otimes \mathbf{1} + \sum_{\Delta^\gamma(t), \Delta^e(s)} (t_{(1)} s_{(1)}^\gamma, s_{(1)}^e) \otimes t_{(2)} \vee s_{(2)} \\ &= (\mathbf{1}, t \vee s) \otimes \mathbf{1} + (\text{Id} \otimes \vee) \Delta^\gamma(t) \Delta^e(s), \end{aligned} \tag{32}$$

where we adopt a modified version of Sweedler's notation

$$\Delta^e s = \sum_{\Delta^e(s)} (s_{(1)}^\gamma, s_{(1)}^e) \otimes s_{(2)},$$

which takes into account the photon component produced by Δ^e . For instance,

$$\begin{aligned}
\Delta^e \Upsilon &= 1 \otimes \Upsilon + (\mathbf{1}, \Upsilon) \otimes \mathbf{1} \\
\Delta^e \check{\Upsilon} &= 1 \otimes \check{\Upsilon} + (\Upsilon, \mathbf{1}) \otimes \Upsilon + (\mathbf{1}, \check{\Upsilon}) \otimes \mathbf{1} \\
\Delta^e \check{\check{\Upsilon}} &= 1 \otimes \check{\check{\Upsilon}} + (\mathbf{1}, \check{\Upsilon}) \otimes \Upsilon + (\mathbf{1}, \check{\check{\Upsilon}}) \otimes \mathbf{1} \\
\Delta^e \check{\check{\check{\Upsilon}}} &= 1 \otimes \check{\check{\check{\Upsilon}}} + 2(\Upsilon, \mathbf{1}) \otimes \check{\Upsilon} + (\check{\Upsilon}, \mathbf{1}) \otimes \Upsilon + (\mathbf{1}, \check{\check{\check{\Upsilon}}}) \otimes \mathbf{1}, \\
\Delta^e \check{\check{\check{\check{\Upsilon}}}} &= 1 \otimes \check{\check{\check{\check{\Upsilon}}}} + (\check{\check{\Upsilon}}, \mathbf{1}) \otimes \Upsilon + (\mathbf{1}, \check{\check{\check{\check{\Upsilon}}}}) \otimes \mathbf{1} \\
\Delta^e \check{\check{\check{\check{\check{\Upsilon}}}}} &= 1 \otimes \check{\check{\check{\check{\check{\Upsilon}}}}} + (\Upsilon, \mathbf{1}) \otimes \check{\check{\Upsilon}} + (\mathbf{1}, \check{\check{\check{\check{\check{\Upsilon}}}}}) \otimes \Upsilon + (\mathbf{1}, \check{\check{\check{\check{\check{\Upsilon}}}}}) \otimes \mathbf{1} \\
\Delta^e \check{\check{\check{\check{\check{\check{\Upsilon}}}}}} &= 1 \otimes \check{\check{\check{\check{\check{\check{\Upsilon}}}}}} + (\Upsilon, \mathbf{1} \otimes \mathbf{1}) \otimes \check{\check{\Upsilon}} + (\mathbf{1}, \check{\check{\Upsilon}}) \otimes \check{\Upsilon} + (\Upsilon, \check{\Upsilon}) \otimes \Upsilon + (\mathbf{1}, \check{\check{\check{\check{\check{\Upsilon}}}}}) \otimes \mathbf{1}, \\
\Delta^e \check{\check{\check{\check{\check{\check{\check{\Upsilon}}}}}}} &= 1 \otimes \check{\check{\check{\check{\check{\check{\check{\Upsilon}}}}}}} + (\mathbf{1}, \check{\check{\Upsilon}}) \otimes \check{\check{\Upsilon}} + (\mathbf{1}, \check{\check{\check{\check{\check{\Upsilon}}}}}) \otimes \Upsilon + (\mathbf{1}, \check{\check{\check{\check{\check{\check{\Upsilon}}}}}}) \otimes \mathbf{1}.
\end{aligned}$$

Let $\epsilon : \mathcal{H}^{\text{qed}} \longrightarrow \mathbb{C}$ be the map which sends everything to 0 except the unit, which is sent to 1.

Proposition 3.4 *The space $\mathcal{H}^{\text{qed}}$ is a bigraded Hopf algebra, not commutative nor cocommutative, of which $\mathcal{H}^\gamma$ is a sub-Hopf algebra.*

Proof. Since all the operations involved in the definition of Δ^{qed} are graded, it is easy to check that the coproduct is a graded operation, that is, it preserves the total degree of each addend. Then, the extended map ϵ is a counit for Δ^{qed} , because each $\Delta^{\text{qed}}(t, s)$ contains exactly the two terms $(t, s) \otimes \mathbf{1}$ and $\mathbf{1} \otimes (t, s)$ in degree $(n, 0)$ and $(0, n)$ respectively, where $n = |t| + |s|$. The proof is by induction. Since by proposition (3.2) it is true on all photon trees, we only need to check it on electron trees, where Δ^{qed} coincides with Δ^e . It is true for $\Delta^e(\mathbf{1})$, so assume that it is true for trees up to order $n - 1$, and consider a tree $t \vee s$ with order n . Then, using (30) and (31) we write

$$\Delta^\gamma(t) = (V(t^r) \otimes \mathbf{1}) \Delta^\gamma(t^l) + F(t),$$

and therefore by (33) we have

$$\begin{aligned}
\Delta^e(t \vee s) &= (\mathbf{1}, t \vee s) \otimes \mathbf{1} + (\text{Id} \otimes \vee)[(V(t^r) \otimes \mathbf{1}) \Delta^\gamma(t^l) + F(t)] \Delta^e(s) \\
&= (\mathbf{1}, t \vee s) \otimes \mathbf{1} + (\text{Id} \otimes \vee)[(V(t^r) \otimes \mathbf{1}) \Delta^\gamma(t^l) \Delta^e(s) + (\text{Id} \otimes \vee) F(t) \Delta^e(s)].
\end{aligned}$$

Because of the presence of the operator $\text{Id} \otimes \vee$, a pair of trees with bidegree $(n, 0)$ can not be contained in the second and third addends, therefore the only such pair is the first addend $t \vee s \otimes \mathbf{1}$. Similarly, a pair of trees with bidegree $(0, n)$ can not be contained in the second addend because of the term $(V(t^r) \otimes \text{Id})$. Therefore it must be contained in the third addend where, by inductive hypothesis, it can only appear as the product of the corresponding terms of bidegrees $(0, |s|)$ and $(0, |t|)$.

Again, the antipode $Sq : \mathcal{H}^{\text{qed}} \longrightarrow \mathcal{H}^{\text{qed}}$ is automatically defined by the recursive formula (29). Since, by definition, Δ^{qed} is a multiplicative map, it only remains to show that it is coassociative. Again, we prove it by induction on the order of trees. Of course, we only need to consider electron trees.

The coassociativity is easy to check for $t = \mathbf{1}$. Suppose that it holds for all trees with order up to n , and let $t \vee s \in Y$ be a tree with order $n + 1$. Then t and s have at most order n , hence, the coassociativity of Δ^{qed} holds for the couple $(t, s) \in Y \otimes Y \subset \mathcal{H}^{\text{qed}}$.

Then, by definition of Δ^{qed} , on one side we have

$$\begin{aligned}
(\Delta^{\text{qed}} \otimes \text{Id}) \Delta^{\text{qed}}(\mathbf{1}, t \vee s) &= (\Delta^{\text{qed}} \otimes \text{Id}) \Delta^e(t \vee s) \\
&= \Delta^e(t \vee s) \otimes \mathbf{1} + \sum_{\Delta^\gamma(t), \Delta^e(s)} \Delta^{\text{qed}}(t_{(1)} s_{(1)}^\gamma, s_{(1)}^e) \otimes t_{(2)} \vee s_{(2)} \\
&= (\mathbf{1}, t \vee s) \otimes 1 \otimes \mathbf{1} + \sum_{\Delta^\gamma(t), \Delta^e(s)} (t_{(1)} s_{(1)}^\gamma, s_{(1)}^e) \otimes (\mathbf{1}, t_{(2)} \vee s_{(2)}) \otimes \mathbf{1} \\
&\quad + \sum_{\Delta^\gamma(t), \Delta^e(s)} \Delta^{\text{qed}}(t_{(1)} s_{(1)}^\gamma, s_{(1)}^e) \otimes t_{(2)} \vee s_{(2)},
\end{aligned}$$

and on the other side

$$\begin{aligned}
(\text{Id} \otimes \Delta^{\text{qed}}) \Delta^{\text{qed}}(\mathbf{1}, t \vee s) &= (\text{Id} \otimes \Delta^{\text{qed}}) \Delta^e(t \vee s) \\
&= (\mathbf{1}, t \vee s) \otimes \mathbf{1} \otimes \mathbf{1} + \sum_{\Delta^\gamma(t), \Delta^e(s)} (t_{(1)}, \mathbf{1})(s_{(1)}^\gamma, s_{(1)}^e) \otimes \Delta^e(t_{(2)} \vee s_{(2)}) \\
&= (\mathbf{1}, t \vee s) \otimes \mathbf{1} \otimes \mathbf{1} + \sum_{\Delta^\gamma(t), \Delta^e(s)} (t_{(1)} \ s_{(1)}^\gamma, s_{(1)}^e) \otimes (\mathbf{1}, t_{(2)} \vee s_{(2)}) \otimes \mathbf{1} \\
&\quad + \sum_{\substack{\Delta^\gamma(t), \Delta^e(s) \\ \Delta^\gamma(t_{(2)}), \Delta^e(s_{(2)})}} (t_{(1)} \ s_{(1)}^\gamma, s_{(1)}^e) \otimes (t_{(21)} \ s_{(21)}^\gamma, s_{(21)}^e) \otimes t_{(22)} \vee s_{(22)}.
\end{aligned}$$

Then, the two sides are equal if and only if

$$\sum_{\Delta^\gamma(t), \Delta^e(s)} \Delta^{\text{qed}}(t_{(1)} \ s_{(1)}^\gamma, s_{(1)}^e) \otimes t_{(2)} \vee s_{(2)} = \sum_{\substack{\Delta^\gamma(t), \Delta^e(s) \\ \Delta^\gamma(t_{(2)}), \Delta^e(s_{(2)})}} (t_{(1)} \ s_{(1)}^\gamma, s_{(1)}^e) \otimes (t_{(21)} \ s_{(21)}^\gamma, s_{(21)}^e) \otimes t_{(22)} \vee s_{(22)}.$$

Since $t \vee s = t' \vee s'$ if and only if $(t, s) = (t', s')$, this equality holds in $\mathcal{H}^{\text{qed}} \otimes \mathcal{H}^{\text{qed}} \otimes \mathcal{H}^e$ if and only if in $\mathcal{H}^{\text{qed}} \otimes \mathcal{H}^{\text{qed}} \otimes \mathcal{H}^{\text{qed}}$ we have

$$\sum_{\Delta^\gamma(t), \Delta^e(s)} \Delta^{\text{qed}}(t_{(1)} \ s_{(1)}^\gamma, s_{(1)}^e) \otimes (t_{(2)}, s_{(2)}) = \sum_{\substack{\Delta^\gamma(t), \Delta^e(s) \\ \Delta^\gamma(t_{(2)}), \Delta^e(s_{(2)})}} (t_{(1)} \ s_{(1)}^\gamma, s_{(1)}^e) \otimes (t_{(21)} \ s_{(21)}^\gamma, s_{(21)}^e) \otimes (t_{(22)}, s_{(22)}).$$

The last equality means that

$$(\Delta^{\text{qed}} \otimes \text{Id}) \Delta^{\text{qed}}(t, s) = (\text{Id} \otimes \Delta^{\text{qed}}) \Delta^{\text{qed}}(t, s),$$

and holds by inductive hypothesis. $\square$

Remark 3.5 On the generators $S_P(t)$ of the algebra $\mathcal{H}^e$, the electron coproduct Δ^e takes the form

$$\Delta^e(S_P(t)) = \sum_{\Delta^e(t)} (t_{(1)}^\gamma, S_P(t_{(1)}^e)) \otimes S_P(t_{(2)}).$$

3.3 Tree expansion for renormalized propagators

Following Ref. [12], and using the fact that the renormalization factors depend only on the vacuum polarization and the self-energy, we know that the photon renormalization factor Z_3 is defined only on the generators $\mathbf{1} \vee t$, and that the electron renormalization factor Z_2 is defined only on elements of the form $S^\gamma(t)$. Therefore, the coefficients $Z_{3,n}$ and $Z_{2,n}$ of the perturbative expansion defined by Eqs. (12) and (13), are defined as

$$Z_{3,n} = \sum_{|t|=n} \zeta_3(V(t)), \tag{33}$$

$$Z_{2,n} = \sum_{|t|=n} \zeta_2(S_P(t)), \tag{34}$$

The maps $\zeta_3 : \mathcal{H}^\gamma \longrightarrow \mathbb{C}$ and $\zeta_2 : \mathcal{H}^e \longrightarrow \mathbb{C}$ are determined recursively on the generators, through a renormalization prescription (cf. [7]),

$$\begin{aligned}
\zeta_3(V(t)) &= \sum_{\pi^\gamma(\Gamma)=V(t)} C_3(\Gamma), \\
\zeta_2(S_P(t)) &= - \sum_{\pi^e(\Gamma)=S_P(t)} C_2(\Gamma),
\end{aligned}$$

and then extended multiplicatively.

Denote by $\zeta : \mathcal{H}^{\text{qed}} \longrightarrow \mathbb{C}$ the product map of ζ_3 and ζ_2 , that is,

$$\zeta(t, s) := \zeta_3(V(t^1)) \dots \zeta_3(V(t^l)) \zeta_2(S_P(s_1)) \dots \zeta_2(S_P(s_k)), \quad (35)$$

where $(t, s) = (V(t^1) \dots V(t^l), S_P(s_1) \dots S_P(s_k))$ is a generic element of $\mathcal{H}^{\text{qed}}$.

In Ref. [7], the renormalized amplitudes corresponding to each tree were found by a recurrence procedure based on the Schwinger equations for the renormalized propagators. In the case of massless QED, it was shown that the relationship between the amplitudes before and after renormalization was encoded by the coproduct on trees. We recall the main results of [7].

Theorem 3.6 *The relation between the coefficients of the expansion (16, 18) of the renormalized and bare photon propagators is*

$$\bar{\varphi}^\gamma(t; q) = \sum_{\Delta^\gamma(t)} \zeta(t_{(1)}, \cdot) \varphi^\gamma(t_{(2)}; q). \quad (36)$$

The relation between the coefficients of the expansion (15, 17) of the renormalized and bare electron propagators is

$$\bar{\varphi}^e(t; q) = \sum_{\Delta^e(t)} \zeta(t_{(1)}^\gamma, t_{(1)}^e) \varphi^e(t_{(2)}; q). \quad (37)$$

In particular, choosing only the generator trees, the same result holds for the renormalization of the vacuum polarization $\Pi(q; e_0)$ and of the self-energy $\Sigma(q; e_0)$.

Theorem 3.7 *The relation between the coefficients of the expansion (20, 21) of the renormalized and bare vacuum polarizations is*

$$\bar{\varphi}^\gamma(V(t); q) = \sum_{\Delta^\gamma(V(t))} \zeta(V(t)_{(1)}, \cdot) \varphi^\gamma(V(t)_{(2)}; q). \quad (38)$$

The relation between the coefficients of the expansion (27, 27) of the renormalized and bare self-energies is

$$\bar{\varphi}^e(S_P(t); q) = \sum_{\Delta^e(t)} \zeta(t_{(1)}^\gamma, S_P(t_{(1)}^e)) \varphi^e(S_P(t_{(2)}); q). \quad (39)$$

4 Renormalization “at the order of interaction”

In section 1, the terms $S_n(q)$ and $D_n(q)$ at each order of interaction are defined using the Feynman’s rules. In fact, they can be regarded as the evaluation of the map U on some formal sums of Feynman’s graphs, that is

$$S_n(q) = U\left(\sum_{\Gamma \in \mathcal{F}^e, |\Gamma|=n} \Gamma\right),$$

$$D_n(q) = U\left(\sum_{\Gamma \in \mathcal{F}^\gamma, |\Gamma|=n} \Gamma\right).$$

Now, the same terms can be regarded as the evaluation of the map $\varphi = \varphi^\gamma \otimes \varphi^e$ on some formal sums of trees

$$S_n(q) = \varphi^e\left(\sum_{|t|=n} t\right) = \langle \varphi, (\cdot, \mathbf{x}_n) \rangle, \quad (40)$$

$$D_n(q) = \varphi^\gamma\left(\sum_{|t|=n} t\right) = \langle \varphi, (\mathbf{u}_n, \cdot) \rangle, \quad (41)$$

where we define the following elements in the Hopf algebra $\mathcal{H}^{\text{qed}}$:

$$\begin{aligned}\mathbf{x}_n &= \sum_{|t|=n} t \in \mathcal{H}^e \\ \mathbf{u}_n &= \sum_{|t|=n} t \in \mathcal{H}^\gamma.\end{aligned}$$

To determine the relationship between the terms of the Green functions at each order of interaction before and after the renormalization, we need to compute the photon and electron coproducts of the elements $\mathbf{x}_n$ and $\mathbf{u}_n$.

4.1 Hopf algebra for photons at order n

Let $\mathbf{u}_n$ and $\mathbf{v}_n$ be the elements of $\mathcal{H}^\gamma$ defined by

$$\mathbf{u}_n = \sum_{|t|=n} t, \quad n \geq 0, \quad (42)$$

$$\mathbf{v}_n = \sum_{|t|=n-1} V(t), \quad n \geq 1. \quad (43)$$

Of course, $\mathbf{v}_n$ is then related to $\mathbf{u}_n$ by the relation $\mathbf{v}_n = V(\mathbf{u}_{n-1})$. Conversely, it is not difficult to show that

$$\mathbf{u}_n = \sum_{k=1}^n P_n^{(k)}(\mathbf{v}_*), \quad (44)$$

where for any $m \geq 1$ and any $k \leq m$, we let

$$P_m^{(k)}(\mathbf{v}_*) = \sum_{\substack{j_1 > 0, \dots, j_k > 0 \\ j_1 + \dots + j_k = m}} \mathbf{v}_{j_1} \cdots \mathbf{v}_{j_k} \quad (45)$$

be the symmetric (noncommutative) polynomial of degree m in k (noncommutative) variables $\mathbf{v}_*$.

Proposition 4.1 *The photon coproduct on trees induces the following coproduct on $\mathbf{v}_n$:*

$$\begin{aligned}\Delta^\gamma \mathbf{v}_1 &= \mathbf{v}_1 \otimes 1 + 1 \otimes \mathbf{v}_1, \\ \Delta^\gamma \mathbf{v}_2 &= \mathbf{v}_2 \otimes 1 + 1 \otimes \mathbf{v}_2, \\ \Delta^\gamma \mathbf{v}_n &= \mathbf{v}_n \otimes 1 + 1 \otimes \mathbf{v}_n + \sum_{m=1}^{n-2} R_m^{n-2}(\mathbf{v}_*) \otimes \mathbf{v}_{n-m}, \quad n \geq 3\end{aligned} \quad (46)$$

where we set

$$R_m^l(\mathbf{v}_*) = \sum_{k=1}^m \binom{l-m+k}{k} P_m^{(k)}(\mathbf{v}_*), \quad m, l \geq 1. \quad (47)$$

A few examples of the values of the polynomials $R_m^n(\mathbf{v}_*)$ might be useful:

$$\begin{aligned}R_1^l(\mathbf{v}_*) &= l\mathbf{v}_1, \\ R_2^l(\mathbf{v}_*) &= (l-1)\mathbf{v}_2 + \frac{l(l-1)}{2}\mathbf{v}_1^2, \\ R_l^l(\mathbf{v}_*) &= \sum_{k=1}^l P_l^{(k)}(\mathbf{v}_*) = \mathbf{u}_l.\end{aligned}$$

Conversely, we can determine each $\mathbf{v}_n$ as a combination of R_m^l 's:

$$\mathbf{v}_n = - \sum_{k=1}^n (-1)^k \binom{n+1}{k+1} R_n^{n+k-1}(\mathbf{v}_*).$$

The computation of $\Delta^\gamma(\mathbf{v}_n)$ for the cases $n = 1$ and $n = 2$ can be done explicitly starting from the coproduct over trees, and using $\Delta^\gamma \Upsilon = \Upsilon \otimes \mathfrak{I} + \mathfrak{I} \otimes \Upsilon$ and $\Delta^\gamma \check{\Upsilon} = \check{\Upsilon} \otimes \mathfrak{I} + \mathfrak{I} \otimes \check{\Upsilon}$. The next terms are:

$$\begin{aligned} \Delta^\gamma \mathbf{v}_3 &= \mathbf{v}_3 \otimes \mathfrak{I} + \mathfrak{I} \otimes \mathbf{v}_3 + \mathbf{v}_1 \otimes \mathbf{v}_2, \\ \Delta^\gamma \mathbf{v}_4 &= \mathbf{v}_4 \otimes \mathfrak{I} + \mathfrak{I} \otimes \mathbf{v}_4 + 2\mathbf{v}_1 \otimes \mathbf{v}_3 + (\mathbf{v}_2 + \mathbf{v}_1^2) \otimes \mathbf{v}_2. \end{aligned}$$

Proposition 4.2 *The photon coproduct evaluated on $\mathbf{u}_n$ gives:*

$$\begin{aligned} \Delta^\gamma \mathbf{u}_1 &= \mathbf{u}_1 \otimes \mathfrak{I} + \mathfrak{I} \otimes \mathbf{u}_1, \\ \Delta^\gamma \mathbf{u}_n &= \mathbf{u}_n \otimes \mathfrak{I} + \mathfrak{I} \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} R_m^n(\mathbf{v}_*) \otimes \mathbf{u}_{n-m}, \quad n \geq 2. \end{aligned} \tag{48}$$

The simplest examples are:

$$\begin{aligned} \Delta^\gamma \mathbf{u}_2 &= \mathbf{u}_1 \otimes \mathfrak{I} + \mathfrak{I} \otimes \mathbf{u}_1 + 2\mathbf{v}_1 \otimes \mathbf{u}_1 \\ \Delta^\gamma \mathbf{u}_3 &= \mathbf{u}_3 \otimes \mathfrak{I} + \mathfrak{I} \otimes \mathbf{u}_3 + 3\mathbf{v}_1 \otimes \mathbf{u}_2 + (2\mathbf{v}_2 + \mathbf{v}_1^2) \otimes \mathbf{u}_1. \end{aligned}$$

Proof of (4.1). The statements in propositions (4.1) and (4.2) are closely related. We prove them by induction simultaneously.

Assume that Eq. (48) is true up to order $n-1$ and that Eq. (46) is true up order n . Then, using Eq. (46) we show that relation (48) is true at order n . Finally, we show that Eq. (46) is true at order $n+1$.

The first part is quite simple. We calculate

$$\begin{aligned} \Delta^\gamma \mathbf{u}_n - \sum_{k=0}^{n-1} (\mathbf{v}_{k+1} \otimes 1) \Delta^\gamma \mathbf{u}_{n-k-1} &= \Delta^\gamma \mathbf{u}_n - \sum_{k=1}^{n-1} (\mathbf{v}_k \otimes 1) \Delta^\gamma \mathbf{u}_{n-k} - \mathbf{v}_n \otimes \mathfrak{I} \\ &= \mathfrak{I} \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} (R_m^n - \mathbf{v}_m - \sum_{k=1}^{m-1} \mathbf{v}_k R_{m-k}^{n-k}) \otimes \mathbf{u}_{n-m} \\ &= \mathfrak{I} \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} R_m^{n-1} \otimes \mathbf{u}_{n-m}. \end{aligned}$$

Introducing this into Eq.(46) yields

$$\begin{aligned} \Delta^\gamma \mathbf{v}_{n+1} &= \mathbf{v}_{n+1} \otimes \mathfrak{I} + (\text{Id} \otimes V)(\mathfrak{I} \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} R_m^{n-1} \otimes \mathbf{u}_{n-m}) \\ &= \mathbf{v}_{n+1} \otimes \mathfrak{I} + \mathfrak{I} \otimes \mathbf{v}_{n+1} + \sum_{m=1}^{n-1} R_m^{n-1} \otimes \mathbf{v}_{n-m+1}. \end{aligned}$$

To derive the last line, we used the fact that $V(\mathbf{u}_n) = \mathbf{v}_{n+1}$.

Knowing $\Delta^\gamma \mathbf{v}_{n+1}$, we can calculate $\Delta^\gamma \mathbf{u}_{n+1}$ through

$$\mathbf{u}_{n+1} = \mathbf{v}_{n+1} + \sum_{k=1}^n \mathbf{v}_{n-k+1} \mathbf{u}_k.$$

We obtain

$$\begin{aligned}
\Delta^\gamma \mathbf{u}_{n+1} &= \Delta^\gamma \mathbf{v}_{n+1} + \sum_{k=1}^n \Delta^\gamma \mathbf{v}_{n-k+1} \Delta^\gamma \mathbf{u}_k \\
&= \mathbf{1} \otimes \mathbf{u}_{n+1} + \mathbf{u}_{n+1} \otimes \mathbf{1} + \sum_{m=1}^n \left(R_m^{n-1} + \mathbf{u}_m + \mathbf{v}_m + \sum_{k=1}^{m-1} \mathbf{v}_k R_{m-k}^{n-k+1} + R_{m-k}^{n-k-1} \mathbf{u}_k \right) \otimes \mathbf{u}_{n+1-m} \\
&\quad + \sum_{l=1}^{n-1} \sum_{i=1}^{n-l} \left(-R_l^{n-1} - \mathbf{u}_l + R_l^{l+i} + R_l^{n-i-1} + \sum_{k=1}^{l-1} -R_k^{n-l+k-1} \mathbf{u}_{l-k} + R_k^{n-l-i+k-1} R_{l-k}^{i+l-k} \right) \otimes \mathbf{v}_{n+1-l-i} \mathbf{u}_i.
\end{aligned}$$

In the above equation, the coefficient of $\mathbf{u}_{n+1-m}$ is denoted by X . It can be simplified by using Eq.(51):

$$X = R_m^{n+1} + \mathbf{u}_m + R_m^{n-1} + \sum_{k=1}^{m-1} R_{m-k}^{n-k-1} \mathbf{u}_k - R_m^n.$$

Equation (57) of the next lemma (4.6), for $l = 0$, together with $R_m^n = \mathbf{u}_m$ gives us $X = R_m^{n+1}$. The coefficient of $\mathbf{v}_{n+1-l-i} \mathbf{u}_i$ is treated similarly. Using Eq.(57), it is shown that the sum of the positive terms is R_l^n and the sum of the negative terms is $-R_l^n$. Therefore, the coefficient of $\mathbf{v}_{n+1-l-i} \mathbf{u}_i$ is zero and the proposition is proved. $\square$

To show lemma (4.6), we shall use several lemmas. The proof of these lemmas is very similar to the proofs given for the non commutative Hopf algebra of formal diffeomorphisms in section 5. Since we think that the results of section 5 can be interesting independently of QED or renormalization theory, the proofs are given with all details in section 5 and not here. We apologize to the reader for this sacrifice of the chronological order.

The first lemma is a relation for $\Delta^\gamma \mathbf{v}_n$ which is deduced from the definition of the coproduct over photon trees.

Lemma 4.3 *For $n \geq 1$, the photon coproduct of $\mathbf{v}_{n+1}$ satisfies the following relation:*

$$\Delta^\gamma \mathbf{v}_{n+1} = \mathbf{v}_{n+1} \otimes 1 + (\text{Id} \otimes V) \left(\Delta^\gamma \mathbf{u}_n - \sum_{k=0}^{n-1} (\mathbf{v}_{k+1} \otimes \mathbf{1}) \Delta^\gamma \mathbf{u}_{n-k-1} \right) \quad (49)$$

Proof. The property described by Eq.(23) together with the fact that $t^l \vee t^r = V(t^r)t^l$, yield a useful identity relating $\mathbf{u}_n$ and $\mathbf{v}_n$:

$$\mathbf{u}_n = \sum_{k=0}^{n-1} \sum_{|t|=k} \sum_{|t^r|=n-k-1} V(t^r) t^l = \sum_{k=0}^{n-1} \mathbf{v}_{n-k} \mathbf{u}_k. \quad (50)$$

From the recursive definition of $\Delta^\gamma V(t)$ (cf. Eq.(28)) and Eq.(23) we find, for $t = t^l \vee t^r$,

$$\begin{aligned}
\Delta^\gamma \mathbf{v}_{n+1} &= \sum_{|t|=n} \Delta^\gamma V(t) = \sum_{|t|=n} V(t) \otimes 1 \\
&\quad + \sum_{|t|=n} (\text{Id} \otimes V) \Delta^\gamma t - \sum_{k=0}^{n-1} (\text{Id} \otimes V) \left(\sum_{|t^r|=k} V(t^r) \otimes \mathbf{1} \right) \sum_{|t^l|=n-k-1} \Delta^\gamma t^l \\
&= \mathbf{v}_{n+1} \otimes 1 + (\text{Id} \otimes V) \Delta^\gamma \mathbf{u}_n - \sum_{k=0}^{n-1} (\text{Id} \otimes V) (\mathbf{v}_{k+1} \otimes \mathbf{1}) \Delta^\gamma \mathbf{u}_{n-k-1}.
\end{aligned}$$

This gives our lemma. $\square$

Then, we need a recursive relation for R_m^n :

Lemma 4.4 *The following recursive relations hold for any $n \geq 1$:*

$$R_m^n(\mathbf{v}_*) = R_m^{n-1}(\mathbf{v}_*) + \mathbf{v}_m + \sum_{l=1}^{m-1} \mathbf{v}_l R_{m-l}^{n-1}(\mathbf{v}_*), \quad m \geq 2 \quad (51)$$

$$R_1^n(\mathbf{v}_*) = R_1^{n-1}(\mathbf{v}_*) + \mathbf{v}_1. \quad (52)$$

Proof. From the obvious identity

$$P_m^{(k)}(\mathbf{v}_*) = \sum_{l=1}^{m-k+1} \mathbf{v}_l P_{m-l}^{(k-1)}(\mathbf{v}_*),$$

we proceed exactly as for the proof of lemma 5.7. We start from the definition

$$R_m^n(\mathbf{v}_*) = \sum_{k=1}^m \binom{n-m+k}{k} P_m^{(k)}(\mathbf{v}_*)$$

and use Pascal's relation for binomial coefficients to obtain the lemma. $\square$

From the recursive relation, we can find the generating function of R_m^n .

Lemma 4.5 *The generating function*

$$R(x, y) = \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} x^n y^{2m} R_m^{n+m}(\mathbf{v}_*) + \frac{1}{1-x} \quad (53)$$

is the Green function of the hamiltonian

$$\mathbf{Z}_3(y) = 1 - \sum_{n=1}^{\infty} y^{2n} \mathbf{v}_n \in \mathcal{H}^\gamma[[y^2]] \quad (54)$$

at energy x . Thus, the following resolvent equation is satisfied

$$R(x, y) = R(z, y) + (x - z) R(x, y) R(z, y). \quad (55)$$

Proof. The proof follows closely the proof of the corresponding lemma (5.9) of section 5. Using the recursive relation (51), we obtain a relation for $R(x, y)$ which leads to the equation

$$R(x, y) = (\mathbf{Z}_3(y) - x)^{-1}, \quad (56)$$

from which the lemma follows. $\square$

It may be noticed that the explicit expression (53) can be interesting for a classical problem of quantum mechanics. Assume that a Hamiltonian H_0 is completely degenerate (i.e. all the eigenvectors of H_0 have the same eigenvalue, that we take equal to 1). What happens to the system when the Hamiltonian is perturbed as $H(y) = H_0 + V(y)$, where $V(y)$ can be expanded $V(y) = \sum_n \mathbf{v}_n y^{2n}$? The answer is given by the Green function $R(x, y) = (H(y) - x)^{-1}$ which can be calculated from Eq.(53).

The final lemma that we need is quite similar to lemma 5.9.

Lemma 4.6 *The following quadratic relations hold for all $l, n \geq 0$:*

$$R_m^{m+n+l+1} = R_m^{m+n} + R_m^{m+l} + \sum_{k=1}^{m-1} R_{m-k}^{m-k+n} R_k^{k+l}, \quad m \geq 2 \quad (57)$$

$$R_1^{n+l} = R_1^n + R_1^l, \quad m = 1. \quad (58)$$

Proof. We now use the resolvent equation in the form

$$\frac{R(x, y)}{1 - z/x} = \frac{R(z, y)}{1 - z/x} + xR(x, y)R(z, y),$$

and follow the proof of lemma 5.9. $\square$

Remark 4.7 As a consequence of the coassociativity of Δ^γ we find

$$\Delta^\gamma R_m^n = R_m^n \otimes 1 + 1 \otimes R_m^n + \sum_{k=1}^{m-1} R_k^n \otimes R_{m-k}^{n-k}.$$

4.2 Hopf structure for electrons at order n

Let $\mathbf{x}_n$ and $\mathbf{y}_n$ be the elements of $\mathcal{H}^e$ defined by

$$\begin{aligned} \mathbf{x}_n &= \sum_{|t|=n} t, \\ \mathbf{y}_n &= \sum_{|t|=n} S_P(t), \end{aligned}$$

for any $n \geq 0$. From the definition of the product antipode S_P over trees, it is straightforward to show that

$$\begin{aligned} \mathbf{y}_n &= S_P(\mathbf{x}_n) = -\mathbf{x}_n - \sum_{m=1}^{n-1} S_P(\mathbf{x}_m)\mathbf{x}_{n-m} = -\mathbf{x}_n - \sum_{m=1}^{n-1} \mathbf{x}_m S_P(\mathbf{x}_{n-m}) \\ &= \sum_{k=1}^n (-1)^k P_n^{(k)}(\mathbf{x}_*). \end{aligned} \quad (59)$$

Notice that, on $\mathbf{x}_n$ and $\mathbf{y}_n$, we have $S_P^2 = \text{Id}$, because the pruning operator on the elements $\mathbf{x}_n$ and $\mathbf{y}_n$ is cocommutative [7]. Hence we also have

$$\mathbf{x}_n = S_P(\mathbf{y}_n) = \sum_{k=1}^n (-1)^k P_n^{(k)}(\mathbf{x}_*).$$

The following proposition describes explicitly the coproduct over $\mathbf{x}_n$.

Proposition 4.8 *The electron coproduct over trees induces the following coproduct on $\mathbf{x}_n \in \mathcal{H}^e$:*

$$\begin{aligned} \Delta^e \mathbf{x}_n &= (1, \mathbf{x}_n) \otimes 1 + 1 \otimes \mathbf{x}_n + \sum_{m=1}^{n-1} T_m^n(\mathbf{v}_*, \mathbf{x}_*) \otimes \mathbf{x}_{n-m}, \\ &= (1, \mathbf{x}_n) \otimes 1 + 1 \otimes \mathbf{x}_n + \sum_{m=1}^{n-1} \bar{T}_m^n(\mathbf{v}_*, \mathbf{y}_*) \otimes \mathbf{x}_{n-m}, \end{aligned} \quad (60)$$

where the polynomials on the left-hand side have mixed terms in both components $\mathcal{H}^\gamma$ and $\mathcal{H}^e$ of the algebra $\mathcal{H}^{\text{qed}}$, defined as

$$\begin{aligned} T_m^n(\mathbf{v}_*, \mathbf{x}_*) &= (R_m^{n-1}(\mathbf{v}_*), 1) + (1, \mathbf{x}_m) + \sum_{k=1}^{m-1} (R_{m-k}^{n-k-1}(\mathbf{v}_*), \mathbf{x}_k), \quad 1 \leq m < n \\ \bar{T}_m^n(\mathbf{v}_*, \mathbf{x}_*) &= (1, \mathbf{x}_m), \end{aligned} \quad (61)$$

and

$$\begin{aligned} \bar{T}_m^n(\mathbf{v}_*, \mathbf{y}_*) &= T_m^n(\mathbf{v}_*, S_P(\mathbf{y}_*)) \\ &= (R_m^{n-1}(\mathbf{v}_*), 1) + \sum_{l=1}^m (-1)^l (1, P_m^{(l)}(\mathbf{y}_*)) + \sum_{k=1}^{m-1} \sum_{l=1}^k (-1)^l (R_{m-k}^{n-k-1}(\mathbf{v}_*), P_k^{(l)}(\mathbf{y}_*)), \end{aligned} \quad (62)$$

Before starting the proof, we give the simplest examples

$$\begin{aligned}\Delta^e \mathbf{x}_1 &= 1 \otimes \mathbf{x}_1 + (\mathbf{1}, \mathbf{x}_1) \otimes \mathbf{1} \\ \Delta^e \mathbf{x}_2 &= 1 \otimes \mathbf{x}_2 + (\mathbf{1}, \mathbf{x}_2) \otimes \mathbf{1} + [(\mathbf{v}_1, \mathbf{1}) + (\mathbf{1}, \mathbf{x}_1)] \otimes \mathbf{x}_1 \\ \Delta^e \mathbf{x}_3 &= 1 \otimes \mathbf{x}_3 + (\mathbf{1}, \mathbf{x}_3) \otimes \mathbf{1} + [(2\mathbf{v}_1, \mathbf{1}) + (\mathbf{1}, \mathbf{x}_1)] \otimes \mathbf{x}_2 + [(\mathbf{v}_2, \mathbf{1}) + (\mathbf{1}, \mathbf{x}_2) + (\mathbf{v}_1, \mathbf{x}_1)] \otimes \mathbf{x}_1.\end{aligned}$$

Proof. The proof will only be sketched, since it follows very closely the proof of the corresponding proposition (4.2) for the photon case.

Equation (60) can be checked by hand up to order 3, by summing the known values of $\Delta^e t$ for $t \leq 3$. We assume that it was proved up to order n , then we start from the recursive definition of the electron coproduct as

$$\Delta^e(t \vee s) = (\mathbf{1}, t \vee s) \otimes \mathbf{1} + (\text{Id} \otimes \vee) \Delta^\gamma t \Delta^e s,$$

to obtain the coproduct of $\mathbf{x}_n$ as

$$\Delta^e \mathbf{x}_{n+1} = (\mathbf{1}, \mathbf{x}_{n+1}) \otimes \mathbf{1} + \sum_{k=0}^n (\text{Id} \otimes \vee) \Delta^\gamma \mathbf{u}_k \Delta^e \mathbf{x}_{n-k}.$$

Then we expand $\Delta^\gamma \mathbf{u}_k$, as given by Eq. (4.2), and $\Delta^e \mathbf{x}_{n-k}$ using Eq. (61) to obtain

$$\begin{aligned}\Delta^e \mathbf{x}_{n+1} &= (\mathbf{1}, \mathbf{x}_{n+1}) \otimes \mathbf{1} + 1 \otimes \mathbf{x}_{n+1} \\ &+ \sum_{m=1}^n \left[(R_m^n, \mathbf{1}) + (\mathbf{1}, \mathbf{x}_m) + \sum_{k=1}^{m-1} R_{m-k}^{n-k}, \mathbf{x}_k \right] \otimes \vee(\mathbf{u}_{n-m}, \mathbf{1}) \\ &+ \sum_{m=1}^{n-1} \left[(\mathbf{1}, T_m^n) + (\mathbf{u}_m, \mathbf{1}) + \sum_{k=1}^{m-1} (\mathbf{1}, \mathbf{u}_k T_{m-k}^{n-k}) \right] \otimes \vee(\mathbf{1}, \mathbf{x}_{n-m}) \\ &+ \sum_{m=1}^{n-1} \sum_{k=m+1}^n \left[(\mathbf{1}, T_m^k) + (R_m^{n-k+m}, \mathbf{1}) + \sum_{l=1}^{m-1} (R_l^{n-k+l}, T_{m-l}^{k-l}) \right] \otimes \vee(\mathbf{u}_{n-k}, \mathbf{x}_{k-m}).\end{aligned}$$

From Eq.(57) and the definition of T_m^k given by Eq.(61), it can be shown that

$$(\mathbf{1}, T_m^k) + (R_m^{n-k+m}, \mathbf{1}) + \sum_{l=1}^{m-1} (R_l^{n-k+l}, T_{m-l}^{k-l}) = (\mathbf{1}, T_m^{n+1}). \quad (63)$$

Moreover, the factor of $\otimes \vee(\mathbf{u}_{n-m}, \mathbf{1})$ is Eq.(63) for $k = m$, and the factor of $\otimes \vee(\mathbf{1}, \mathbf{x}_{n-m})$ is Eq.(63) for $k = n$. Therefore,

$$\begin{aligned}\Delta^e \mathbf{x}_{n+1} &= (\mathbf{1}, \mathbf{x}_{n+1}) \otimes \mathbf{1} + 1 \otimes \mathbf{x}_{n+1} + (\mathbf{1}, T_n^{n+1}) \otimes \mathbf{x}_1 \\ &+ \sum_{m=1}^{n-1} (\mathbf{1}, T_m^{n+1}) \otimes \left(\sum_{k=m}^n \vee(\mathbf{u}_{n-k}, \mathbf{x}_{k-m}) \right) \\ &= (\mathbf{1}, \mathbf{x}_{n+1}) \otimes \mathbf{1} + 1 \otimes \mathbf{x}_{n+1} + \sum_{m=1}^n (\mathbf{1}, T_m^{n+1}) \otimes \mathbf{x}_{n+1-m}.\end{aligned}$$

The last line was obtained because

$$\begin{aligned}\sum_{k=m}^n \vee(\mathbf{u}_{n-k}, \mathbf{x}_{k-m}) &= \sum_{k=0}^{n-m} \vee(\mathbf{u}_k, \mathbf{x}_{n-m-k}) \\ &= \sum_{k=0}^{n-m} \sum_{|s|=k} \sum_{|t|=n-m-k} \vee(s, t) \\ &= \sum_{k=0}^{n-m} \sum_{|s|=k} \sum_{|t|=n-m-k} s \vee t = \sum_{|t|=n-m+1} t = \mathbf{x}_{n-m+1}.\end{aligned}$$

□

From the expression for $T_m^n(\mathbf{v}_*, \mathbf{x}_*)$ given in Eq. (61), and that for the photon generating function $R(x, y)$ given in Eq. (56), it is straightforward to show the following lemma.

Lemma 4.9 *The generating function for the terms $T_m^n(\mathbf{v}_*, \mathbf{x}_*)$ is*

$$T(x, y) = \sum_{m \geq 1} \sum_{n \geq 0} x^n y^{2m} T_m^{m+n}(\mathbf{v}_*, \mathbf{x}_*) + \frac{1}{1-x}$$

and satisfies the equation

$$T(x, y) = (\mathbf{Z}_3(y)(\mathbf{Z}_3(y) - x)^{-1}, \mathbf{Z}_2(y)),$$

where $\mathbf{Z}_3(y)$ was defined in (54) and

$$\mathbf{Z}_2(y) = 1 + \sum_{m \geq 1} y^{2m} \mathbf{x}_m \in \mathcal{H}^e[[y^2]].$$

We can now determine the action of the electron coproduct Δ^e on the elements $\mathbf{y}_n = S_P(\mathbf{x}_n)$.

Proposition 4.10 *The product antipode intertwines with the electron coproduct according to the formula*

$$\Delta^e S_P = ((\text{Id}, S_P) \otimes S_P) \Delta^e. \quad (64)$$

In other words, the electron coproduct of $\mathbf{y}_n$ is given by

$$\Delta^e \mathbf{y}_n = (\mathbf{1}, \mathbf{y}_n) \otimes \mathbf{1} + 1 \otimes \mathbf{y}_n + \sum_{m=1}^{n-1} T_m^n(\mathbf{v}_*, \mathbf{y}_*) \otimes \mathbf{y}_{n-m}. \quad (65)$$

Proof. The proof of Eq.(65) is recursive and follows the now usual scheme. The electron coproduct is applied to Eq.(59), and gives

$$\Delta^e \mathbf{y}_n = -\Delta^e \mathbf{x}_n - \sum_{m=1}^{n-1} \Delta^e \mathbf{y}_m \Delta^e \mathbf{x}_{n-m}.$$

We expand all coproducts on the right-hand side. The expansion yields a term $1 \otimes \mathbf{y}_n + (\mathbf{1}, \mathbf{y}_n) \otimes \mathbf{1}$, plus a term in $\otimes \mathbf{x}_{n-m}$, a term in $\otimes \mathbf{y}_{n-m}$ and a term in $\otimes \mathbf{y}_{k-m} \mathbf{x}_{n-k}$. From Eq.(59) and Eq.(57), it turns out that the coefficient of $\mathbf{x}_{n-m}$ is $-(R_m^{n-1}, \mathbf{1})$, the coefficient of $\mathbf{y}_{k-m} \mathbf{x}_{n-k}$ is also $-(R_m^{n-1}, \mathbf{1})$, and the coefficient of $\mathbf{y}_{n-m}$ is $(\mathbf{1}, \mathbf{y}_m) + \sum_k (R_{m-k}^{n-k-1}, \mathbf{y}_k)$. Using again Eq.(59) we obtain

$$\Delta^e \mathbf{y}_n = (\mathbf{1}, \mathbf{y}_n) \otimes \mathbf{1} + 1 \otimes \mathbf{y}_n + \sum_{m=1}^{n-1} [(R_m^{n-1}, \mathbf{1}) + (\mathbf{1}, \mathbf{y}_m) + \sum_{k=1}^{m-1} (R_{m-k}^{n-k-1}, \mathbf{y}_k)] \otimes \mathbf{y}_{n-m},$$

which is Eq.(65) at order n . □

The elements $\mathbf{x}_n$ are used to calculate the electron propagator and the elements $\mathbf{y}_n$ are used to calculate the electron self-energy.

4.3 Renormalization at order n

In ref.[7], the renormalized amplitudes were defined recursively on each tree. However, they are *not* a finite quantity. As shown in Ref.[7], finite renormalized amplitudes are obtained by summing all trees of a given order. Applying the results of section 3.3 and the formulas of propositions (4.2), (4.1) and (4.8), (4.10) for the coproducts, we can describe the renormalization of the propagators at each order of interaction.

The coefficients $Z_{3,n}$ and $Z_{2,n}$ of the perturbative expansion defined by Eqs. (12) and (13), are now expanded over the elements $\mathbf{v}_*$ and $\mathbf{y}_*$ as

$$Z_{3,n} = \sum_{|t|=n} \zeta_3(V(t)) = \zeta_3(\mathbf{v}_n), \quad (66)$$

$$Z_{2,n} = \sum_{|t|=n} \zeta_2(S_P(t)) = \zeta_2(\mathbf{y}_n), \quad (67)$$

where the multiplicative maps $\zeta_3 : \mathcal{H}^\gamma \longrightarrow \mathbb{C}$ and $\zeta_2 : \mathcal{H}^e \longrightarrow \mathbb{C}$ were defined in section 3.3.

Theorem 4.11 *The photon propagator at each order n of interaction is renormalized as*

$$\bar{D}_n(q) = D_n(q) + \sum_{m=1}^n R_m^n(Z_{3,*}) D_{n-m}(q),$$

where the noncommutative polynomial R_m^n was defined in (47). Similarly, the electron propagator at each order n of interaction is renormalized as

$$\bar{S}_n(q) = S_n(q) + \sum_{m=1}^n \bar{T}_m^n(Z_{3,*}, Z_{2,*}) S_{n-m}(q),$$

where the noncommutative polynomial T_m^n was defined in (62).

Proof. Start with the definitions

$$\begin{aligned} D_n(q) &= \varphi^\gamma(\mathbf{u}_n), & \bar{D}_n(q) &= \bar{\varphi}^\gamma(\mathbf{u}_n), \\ S_n(q) &= \varphi^e(\mathbf{x}_n), & \bar{S}_n(q) &= \bar{\varphi}^e(\mathbf{x}_n), \end{aligned}$$

with renormalized amplitudes

$$\begin{aligned} \bar{\varphi}^\gamma &= \zeta_3 \star \varphi^\gamma, \\ \bar{\varphi}^e &= (\zeta_3, \zeta_2) \star \varphi^e. \end{aligned}$$

Then, for the photon propagator we have:

$$\begin{aligned} \bar{D}_n(q) &= \bar{\varphi}^\gamma(\mathbf{u}_n) \\ &= (\zeta_3 \star \varphi^\gamma)(\mathbf{u}_n) \\ &= \zeta_3(\mathbf{u}_n) \varphi^\gamma(\mathbf{1}) + \zeta_3(\mathbf{1}) \varphi^\gamma(\mathbf{u}_n) + \sum_{m=1}^{n-1} R_m^n(\zeta_3(\mathbf{v}_*)) \varphi^\gamma(\mathbf{u}_{n-m}) \\ &= \zeta_3(R_n^n(\mathbf{v}_*)) \varphi^\gamma(\mathbf{1}) + \zeta_3(\mathbf{1}) \varphi^\gamma(\mathbf{u}_n) + \sum_{m=1}^{n-1} R_m^n(\zeta_3(\mathbf{v}_*)) \varphi^\gamma(\mathbf{u}_{n-m}) \\ &= D_n(q) + \sum_{m=1}^n R_m^n(Z_{3,*}) D_{n-m}(q). \end{aligned}$$

A similar computation yields the renormalization of the electron propagator. □

Theorem 4.12 *The vacuum polarization at each order n of interaction is renormalized as*

$$\bar{\Pi}_n(q) = \Pi_n(q) + Z_{3,n} + \sum_{m=1}^{n-2} R_m^{n-2}(Z_{3,*}) \Pi_{n-m}(q),$$

where the noncommutative polynomial R_m^n was defined in (47). Similarly, the self-energy at each order n of interaction is renormalized as

$$\bar{\Sigma}_n(q) = \Sigma_n(q) - Z_{2,n} + \sum_{m=1}^{n-1} T_m^n(Z_{3,*}, Z_{2,*}) \Sigma_{n-m}(q),$$

where the noncommutative polynomial T_m^n was defined in (61).

Proof. Start with the definitions

$$\begin{aligned}\Pi_n(q) &= \varphi^\gamma(\mathbf{v}_n), & \bar{\Pi}_n(q) &= \bar{\varphi}^\gamma(\mathbf{v}_n), \\ \Sigma_n(q) &= -\varphi^e(\mathbf{y}_n), & \bar{\Sigma}_n(q) &= -\bar{\varphi}^e(\mathbf{y}_n),\end{aligned}$$

with renormalized amplitudes as above. Then, for the vacuum polarization we have:

$$\begin{aligned}\bar{\Pi}_n(q) &= \bar{\varphi}^\gamma(\mathbf{v}_n) \\ &= (\zeta_3 \star \varphi^\gamma)(\mathbf{v}_n) \\ &= \zeta_3(\mathbf{v}_n)\varphi^\gamma(\mathbf{1}) + \zeta_3(\mathbf{1})\varphi^\gamma(\mathbf{v}_n) + \sum_{m=1}^{n-2} R_m^{n-2}(\zeta_3(\mathbf{v}_*))\varphi^\gamma(\mathbf{v}_{n-m}) \\ &= Z_{3,n} + \Pi_n(q) + \sum_{m=1}^{n-2} R_m^{n-2}(Z_{3,*})\Pi_{n-m}(q).\end{aligned}$$

A similar computation gives the formula for the renormalization of the self-energy. $\square$

5 Noncommutative Hopf algebra of formal diffeomorphism

It is well known that the composition of series with non-commutative coefficients is not an associative operation. However, in quantum field theory, the compositions of matrix-valued series are necessary, and the associativity of the operations is usually imposed by replacing the usual composition with a “normal ordering”, which re-establishes the position of the misplaced variables. Such an operation is not binary, and needs to be defined “a priori” on each set of n variables.

In this section, we define a dual operation to the normal ordering, which surprisingly is a binary co-operation, that is, a coproduct.

In fact, it turns out that this coproduct is part of the structure of a non-commutative nor commutative Hopf algebra which plays the role of “dual” to the space of formal diffeomorphisms $\mathbf{f}(x) = x + \sum_n x^{n+1}\mathbf{a}_n$ where $\mathbf{a}_n$ belongs to a unital associative (non-commutative) algebra and x commutes with all $\mathbf{a}_n$ ’s. That is, its abelianization is the commutative Hopf algebra of functions on formal diffeomorphisms of $\mathbb{C}$.

As the group of formal diffeomorphisms acts on the set of formal power series, we finally find a “dual” non-commutative version given by a coaction of the Hopf algebra on the free associative algebra on infinitely many variables. This coaction will be used to compose non-commutative formal series and non-commutative formal diffeomorphisms in normal ordering.

5.1 The Hopf algebra

Definition 5.1 Let $\mathcal{H}^{\text{dif}} = \mathbb{C}\langle \mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3 \dots \rangle$ be the algebra of noncommutative polynomials on infinitely many variables $\mathbf{a}_n$, for $n \geq 1$, with unit 1. For any $m \geq 1$ and any $k \leq m$, let

$$P_m^{(k)}(\mathbf{a}_*) = \sum_{\substack{j_1 > 0, \dots, j_k > 0 \\ j_1 + \dots + j_k = m}} \mathbf{a}_{j_1} \cdots \mathbf{a}_{j_k} \quad (68)$$

be the symmetric (noncommutative) polynomial of degree m in k variables $\mathbf{a}_*$. For instance,

$$\begin{aligned}P_1^{(1)}(\mathbf{a}_*) &= \mathbf{a}_1, \\ P_2^{(1)}(\mathbf{a}_*) &= \mathbf{a}_2, & P_2^{(2)}(\mathbf{a}_*) &= \mathbf{a}_1^2, \\ P_3^{(1)}(\mathbf{a}_*) &= \mathbf{a}_3, & P_3^{(2)}(\mathbf{a}_*) &= \mathbf{a}_2\mathbf{a}_1 + \mathbf{a}_1\mathbf{a}_2, & P_3^{(3)}(\mathbf{a}_*) &= \mathbf{a}_1^3.\end{aligned}$$

Proposition 5.2 *The algebra $\mathcal{H}^{\text{dif}}$ is a Hopf algebra with coproduct defined on the generators as*

$$\Delta^{\text{dif}} \mathbf{a}_n = 1 \otimes \mathbf{a}_n + \mathbf{a}_n \otimes 1 + \sum_{m=1}^{n-1} Q_m^{n-m}(\mathbf{a}_*) \otimes \mathbf{a}_{n-m}, \quad (69)$$

where we set

$$Q_m^l(\mathbf{a}_*) = \sum_{k=1}^m \binom{l+1}{k} P_m^{(k)}(\mathbf{a}_*), \quad l \geq 0, \quad m \geq 1, \quad (70)$$

and where we assume that $\binom{l+1}{k}$ is zero for $k > l+1$. The counit is given by $\epsilon(\mathbf{a}_n) = 0$ and $\epsilon(1) = 1$, and the antipode is given by the standard recursive formula

$$\begin{aligned} S^{\text{dif}} \mathbf{a}_n &= -\mathbf{a}_n - \sum_{m=1}^{n-1} S^{\text{dif}}(Q_m^{n-m}(\mathbf{a}_*)) \mathbf{a}_{n-m} \\ &= -\mathbf{a}_n - \sum_{m=1}^{n-1} Q_m^{n-m}(\mathbf{a}_*) S^{\text{dif}} \mathbf{a}_{n-m}. \end{aligned}$$

Here are some examples of coproducts:

$$\begin{aligned} \Delta^{\text{dif}} \mathbf{a}_1 &= \mathbf{a}_1 \otimes 1 + 1 \otimes \mathbf{a}_1, \\ \Delta^{\text{dif}} \mathbf{a}_2 &= \mathbf{a}_2 \otimes 1 + 1 \otimes \mathbf{a}_2 + 2\mathbf{a}_1 \otimes \mathbf{a}_1, \\ \Delta^{\text{dif}} \mathbf{a}_3 &= \mathbf{a}_3 \otimes 1 + 1 \otimes \mathbf{a}_3 + 3\mathbf{a}_1 \otimes \mathbf{a}_2 + 2\mathbf{a}_2 \otimes \mathbf{a}_1 + \mathbf{a}_1^2 \otimes \mathbf{a}_1, \end{aligned}$$

and some examples of antipodes:

$$\begin{aligned} S^{\text{dif}} \mathbf{a}_1 &= -\mathbf{a}_1, \\ S^{\text{dif}} \mathbf{a}_2 &= -\mathbf{a}_2 + 2\mathbf{a}_1^2, \\ S^{\text{dif}} \mathbf{a}_3 &= -\mathbf{a}_3 + 3\mathbf{a}_1 \mathbf{a}_2 + 2\mathbf{a}_2 \mathbf{a}_1 - 5\mathbf{a}_1^3. \end{aligned}$$

The coproduct is clearly not cocommutative, and the square of the antipode is equal to the identity only for $\mathbf{a}_1$ and $\mathbf{a}_2$.

The proof of the proposition is given after a few remarks.

Corollary 5.3 *Let $\mathcal{H}^{\text{odd}} = \mathcal{H}^{\text{dif}}/(\mathbf{a}_{2n+1}, n \geq 0)$ be the quotient algebra of $\mathcal{H}^{\text{dif}}$ obtained by setting all the odd coefficients to zero. Then $\mathcal{H}^{\text{odd}}$ is a Hopf algebra with coproduct*

$$\Delta^{\text{odd}} \mathbf{a}_{2n} = 1 \otimes \mathbf{a}_{2n} + \mathbf{a}_{2n} \otimes 1 + \sum_{m=1}^{n-1} \bar{Q}_m^{n-m}(\mathbf{a}_{2*}) \otimes \mathbf{a}_{2(n-m)}, \quad (71)$$

where we set

$$\bar{Q}_m^l(\mathbf{a}_{2*}) = Q_m^{2l}(\mathbf{a}_{2*}) = \sum_{k=1}^m \binom{2l+1}{k} \bar{P}_m^{(k)}(\mathbf{a}_{2*}), \quad l \geq 0, \quad m \geq 1, \quad (72)$$

and

$$\bar{P}_m^{(k)}(\mathbf{a}_{2*}) = \sum_{\substack{j_1 > 0, \dots, j_k > 0 \\ j_1 + \dots + j_k = m}} \mathbf{a}_{2j_1} \cdots \mathbf{a}_{2j_k}.$$

Remark 5.4 The coproduct Δ^{dif} on $\mathcal{H}^{\text{dif}}$ induces a Hopf structure on the commutative algebra $\mathcal{H}_{ab}^{\text{dif}} = \mathcal{H}^{\text{dif}}/([\mathcal{H}^{\text{dif}}, \mathcal{H}^{\text{dif}}]) \cong \mathbb{C}[a_1, a_2, a_3, \dots]$. Since on commutative variables the polynomial $P_m^{(k)}(a_*)$ becomes

$$P_m^{(k)}(a_*) = \sum_{\substack{p_1 + 2p_2 + \dots + mp_m = m \\ p_1 + p_2 + \dots + p_m = k}} \frac{k!}{p_1! \cdots p_m!} a_1^{p_1} \cdots a_m^{p_m},$$

where the sum runs over $p_1, \dots, p_m \geq 0$, the coproduct on commutative variables becomes

$$\begin{aligned} \Delta^{\text{dif}} a_n &= 1 \otimes a_n + a_n \otimes 1 \\ &+ \sum_{m=1}^{n-1} \sum_{k=1}^m \frac{(n-m+1)!}{(n-m+1-k)!} \sum_{\substack{p_1+\dots+m p_m=m \\ p_1+p_2+\dots+p_m=k}} \frac{1}{p_1! \dots p_m!} a_1^{p_1} \dots a_m^{p_m} \otimes a_{n-m}. \end{aligned}$$

Remark 5.5 The commutative Hopf algebra $\mathcal{H}_{ab}^{\text{dif}}$ is the dual Hopf algebra of the group

$$\mathcal{G}^{\text{dif}} = \left\{ f : \mathbb{C} \longrightarrow \mathbb{C}, f(x) = x + \sum_{n \geq 1} x^{n+1} a_n(f), a_n \in \mathbb{C} \right\}$$

of formal diffeomorphisms f of $\mathbb{C}$ such that $f(0) = 0$ and $f'(0) = 1$. Hence $\mathcal{H}^{\text{dif}}$ can be considered as a non commutative analogue of (the commutative part of) the Connes-Moscovici Hopf algebra used in [9], [10], [11].

To see this, we show that the generators a_n of $\mathcal{H}_{ab}^{\text{dif}}$ behave as the functions on $\mathcal{G}^{\text{dif}}$ defined as

$$\langle a_n, f \rangle = a_n(f) = \frac{1}{(n+1)!} \frac{d^{n+1} f(0)}{dx^{n+1}}.$$

That is, we show that the composition $f \circ g$ of two series $f, g \in \mathcal{G}^{\text{dif}}$ gives rise to the coproduct Δ^{dif} by the duality $\langle \Delta^{\text{dif}} a_n, g \otimes f \rangle = \langle a_n, f \circ g \rangle$, where the pairing on the left hand-side is defined as $\langle a_n \otimes a_m, g \otimes f \rangle = a_n(g) a_m(f)$.

Let f and g be two series in $\mathcal{G}^{\text{dif}}$, then

$$\begin{aligned} f \circ g(x) &= g(x) + \sum_{n=1}^{\infty} g(x)^{n+1} a_n(f) \\ &= x + \sum_{n=1}^{\infty} x^{n+1} a_n(g) + \sum_{n=1}^{\infty} x^{n+1} (1 + \sum_{j=1}^{\infty} x^j a_j(g))^{n+1} a_n(f) \\ &= x + \sum_{n=1}^{\infty} x^{n+1} a_n(g) + \sum_{n=1}^{\infty} x^{n+1} \sum_{k=0}^{n+1} \binom{n+1}{k} \left(\sum_{j=1}^{\infty} x^j a_j(g) \right)^k a_n(f) \\ &= x + \sum_{n=1}^{\infty} x^{n+1} (a_n(g) + a_n(f)) + \sum_{n=1}^{\infty} \sum_{k=1}^{n+1} \sum_{m=k}^{\infty} x^{n+m+1} \binom{n+1}{k} P_m^{(k)}(a_*(g)) a_n(f). \end{aligned}$$

If we identify the terms of order x^{n+1} we obtain precisely

$$a_n(f \circ g) = a_n(f) + a_n(g) + \sum_{m=1}^{n-1} \sum_{k=1}^m \binom{n-m+1}{k} P_m^{(k)}(a_*(g)) a_{n-m}(f).$$

Remark 5.6 Similarly to (5.4) and (5.5), the abelianized Hopf algebra $\mathcal{H}_{ab}^{\text{odd}} = \mathcal{H}^{\text{dif}} / ([\mathcal{H}^{\text{dif}}, \mathcal{H}^{\text{dif}}]) \cong \mathbb{C}[a_2, a_4, a_6, \dots]$, with the coproduct induced by Δ^{odd} , is dual to the group

$$\mathcal{G}^{\text{odd}} = \left\{ f : \mathbb{C} \longrightarrow \mathbb{C}, f(x) = x + \sum_{n \geq 1} x^{2n+1} a_{2n}(f), a_{2n} \in \mathbb{C} \right\}$$

of “odd” formal diffeomorphisms considered in [12].

Proof of (5.2). The only non trivial statement is the coassociativity of the coproduct Δ^{dif} . Applying the definition (69), we obtain

$$\begin{aligned} (\Delta^{\text{dif}} \otimes 1) \Delta^{\text{dif}} \mathbf{a}_n - (1 \otimes \Delta^{\text{dif}}) \Delta^{\text{dif}} \mathbf{a}_n &= \\ &= \sum_{m=1}^{n-1} \left(\Delta^{\text{dif}} Q_m^{n-m} - 1 \otimes Q_m^{n-m} - Q_m^{n-m} \otimes 1 - \sum_{k=1}^{m-1} Q_k^{n-k} \otimes Q_{m-k}^{n-m} \right) \otimes \mathbf{a}_{n-m}. \end{aligned}$$

The right-hand side is zero because of lemma (5.10), hence the coproduct is coassociative. $\square$

We prove lemma (5.10) in several steps, starting with a recursion formula for Q_m^n . First, let us give some examples of the polynomials Q_m^n . We have $Q_m^0 = \mathbf{a}_m$ for any $m \geq 1$, and

$$\begin{aligned} Q_1^l &= (l+1)\mathbf{a}_1, \\ Q_2^l &= (l+1)\mathbf{a}_2 + \frac{l(l+1)}{2}\mathbf{a}_1^2, \\ Q_3^l &= (l+1)\mathbf{a}_3 + \frac{l(l+1)}{2}(\mathbf{a}_1\mathbf{a}_2 + \mathbf{a}_2\mathbf{a}_1) + \frac{(l+1)l(l-1)}{6}\mathbf{a}_1^3 \end{aligned}$$

for any $l \geq 1$.

Lemma 5.7 *The following recursive identities are satisfied:*

$$\begin{aligned} Q_m^n &= Q_m^{n-1} + \mathbf{a}_m + \sum_{l=1}^{m-1} \mathbf{a}_l Q_{m-l}^{n-1}, \quad n \geq 1, m \geq 2 \\ Q_1^n &= Q_1^{n-1} + \mathbf{a}_1, \quad n \geq 1, m = 1. \end{aligned} \tag{73}$$

Proof. From the definition of $P_m^{(k)}$ it is clear that, for $k > 1$,

$$P_m^{(k)} = \sum_{l=1}^{m+1-k} \mathbf{a}_l P_{m-l}^{(k-1)}.$$

Therefore,

$$\begin{aligned} Q_m^n &= \sum_{k=1}^m \binom{n+1}{k} P_m^{(k)} = \sum_{k=1}^m \binom{n}{k} P_m^{(k)} + \sum_{k=1}^m \binom{n}{k-1} P_m^{(k)} \\ &= Q_m^{n-1} + \mathbf{a}_m + \sum_{k=1}^{m-1} \binom{n}{k} P_m^{(k+1)} \\ &= Q_m^{n-1} + \mathbf{a}_m + \sum_{k=1}^{m-1} \sum_{l=1}^{m-k} \binom{n}{k} \mathbf{a}_l P_{m-l}^{(k)} \\ &= Q_m^{n-1} + \mathbf{a}_m + \sum_{l=1}^{m-1} \mathbf{a}_l Q_{m-l}^{n-1}. \end{aligned}$$

The identity $Q_1^n = Q_1^{n-1} + \mathbf{a}_1$ comes from the definition of $Q_1^n = (n+1)\mathbf{a}_1$. $\square$

Lemma 5.8 *The generating function*

$$Q(x, y) = \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} x^n y^m Q_m^n(\mathbf{a}_*) + \frac{1}{1-x} \tag{74}$$

is the Green function of the Hamiltonian $y/\mathbf{f}(y)$ at x , where

$$\mathbf{f}(y) = y + \sum_{n=1}^{\infty} y^{n+1} \mathbf{a}_n \in \mathcal{H}^{\text{dif}}[[y]].$$

That is, the generating function is

$$Q(x, y) = (y - x\mathbf{f}(y))^{-1} \mathbf{f}(y) = (y\mathbf{f}(y)^{-1} - x)^{-1}, \tag{75}$$

and satisfies the following resolvent equation

$$Q(x, y) = Q(z, y) + (x - z)Q(x, y)Q(z, y). \tag{76}$$

Proof. Using lemma 5.7, we can write $Q(x, y)$ as

$$\begin{aligned} Q(x, y) &= \frac{1}{1-x} + \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} x^n y^m Q_m^{n-1} + \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} y^m x^n \mathbf{a}_m \\ &\quad + \sum_{m=2}^{\infty} \sum_{n=1}^{\infty} \sum_{l=1}^{m-1} x^n y^m \mathbf{a}_l Q_{m-l}^{n-1}. \end{aligned}$$

Setting $\mathbf{f}(y) = y + \sum_{n=1}^{\infty} y^{n+1} \mathbf{a}_n$, we can rewrite the above equation as

$$\begin{aligned} Q(x, y) &= \frac{1}{1-x} + x \left(Q(x, y) - \frac{1}{1-x} \right) + \left(\frac{\mathbf{f}(y)}{y} - 1 \right) \frac{1}{1-x} \\ &\quad + x \left(\frac{\mathbf{f}(y)}{y} - 1 \right) \left(Q(x, y) - \frac{1}{1-x} \right), \end{aligned}$$

and obtain the following identity

$$[y - x\mathbf{f}(y)]Q(x, y) = \mathbf{f}(y), \quad (77)$$

from which follows Eq. (75).

Now, multiply the left hand-side of Eq. (76) by $[y - x\mathbf{f}(y)]$ from the left, and using Eq. (77), we obtain:

$$\begin{aligned} [y - x\mathbf{f}(y)][Q(z, y) + (x - z)Q(x, y)Q(z, y)] &= \\ &= [y - x\mathbf{f}(y)]Q(z, y) + (x - z)[y - x\mathbf{f}(y)]Q(x, y)Q(z, y) \\ &= [y - z\mathbf{f}(y)]Q(z, y) + (z - x)\mathbf{f}(y)Q(z, y) + (x - z)\mathbf{f}(y)Q(z, y) \\ &= \mathbf{f}(y) = [y - x\mathbf{f}(y)]Q(x, y). \end{aligned}$$

This proves the resolvent equation (76). $\square$

Lemma 5.9 *The following quadratic equations are satisfied for all $l, n \geq 0$:*

$$\begin{aligned} Q_m^{n+l+1} &= Q_m^n + Q_m^l + \sum_{k=1}^{m-1} Q_k^l Q_{m-k}^n, \quad m \geq 2, \\ Q_1^{n+l+1} &= Q_1^n + Q_1^l. \end{aligned} \quad (78)$$

Proof. To prove this equations, we rewrite the resolvent equation (76) as

$$\frac{Q(z, y)}{1 - x/z} = \frac{Q(x, y)}{1 - x/z} + zQ(x, y)Q(z, y).$$

On the left-hand side we have

$$\frac{Q(z, y)}{1 - x/z} = \frac{z}{(z - x)(1 - z)} + \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} y^m x^l z^{n-l} Q_m^n,$$

while on the right-hand side the first term is

$$\frac{Q(x, y)}{1 - x/z} = \frac{z}{(z - x)(1 - x)} + \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} y^m x^{n+l} z^{-l} Q_m^n,$$

and from definition (74) we obtain the second term

$$\begin{aligned} zQ(x, y)Q(z, y) &= \sum_{m=2}^{\infty} \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \left(\sum_{k=1}^{m-1} y^m x^l z^{n+1} Q_k^l Q_{m-k}^n \right) \\ &\quad + \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} y^m x^l z^{n+1} (Q_m^l + Q_m^n) + \frac{z}{(1 - x)(1 - z)}. \end{aligned}$$

The fractions cancel out and the coefficient of $y^m x^l z^{n+1}$ ($m > 0, l \geq 0, n \geq 0$) gives Eq.(78). $\square$

Lemma 5.10 *The coproduct of $Q_m^n(\mathbf{a}_*)$ is given by*

$$\begin{aligned}\Delta^{\text{dif}} Q_m^n(\mathbf{a}_*) &= 1 \otimes Q_m^n + Q_m^n \otimes 1 + \sum_{k=1}^{m-1} Q_k^{n+m-k} \otimes Q_{m-k}^n, \quad m \geq 2, n \geq 0 \\ \Delta^{\text{dif}} Q_1^n(\mathbf{a}_*) &= 1 \otimes Q_1^n + Q_1^n \otimes 1, \quad m = 1, n \geq 0.\end{aligned}\tag{79}$$

Proof. The proof is given by a recursion argument. From the definition of the coproduct given in Eq. (69), the identity (79) is true for $n = 0$. Now assume that relation (79) holds up to order n . Then, using Eq.(73), we have

$$\Delta^{\text{dif}} Q_m^{n+1} = \Delta^{\text{dif}} Q_m^n + \Delta^{\text{dif}} \mathbf{a}_m + \sum_{l=1}^{m-1} \Delta^{\text{dif}} \mathbf{a}_l \Delta^{\text{dif}} Q_{m-l}^n.$$

Expanding these coproducts we obtain

$$\begin{aligned}\Delta^{\text{dif}} Q_m^{n+1} &= 1 \otimes Q_m^{n+1} + Q_m^{n+1} \otimes 1 + \sum_{k=1}^{m-1} \left(Q_k^{m-k} + Q_k^n + \sum_{p=1}^{k-1} Q_p^{m-k} Q_{k-p}^n \right) \otimes \mathbf{a}_{m-k} \\ &\quad + \sum_{k=1}^{m-1} \left(Q_k^{n+m-k} + \mathbf{a}_k + \sum_{l=1}^{k-1} \mathbf{a}_l Q_{k-l}^{n+m-k} \right) \otimes Q_{m-k}^n \\ &\quad + \sum_{k=1}^{m-1} \sum_{l=1}^{m-k-1} \left(Q_k^{n+m-k-l} + Q_k^l + \sum_{p=1}^{k-1} Q_p^l Q_{k-p}^{n+m-k-l} \right) \otimes \mathbf{a}_l Q_{m-k-l}^n.\end{aligned}$$

From Eq.(78), we see that the each sum in parenthesis is equal to $Q_k^{n+1+m-k}$. Therefore, using Eq.(73), we obtain Eq.(79) for $n + 1$. $\square$

5.2 Coaction on the dual of series

Proposition 5.11 *Let $\mathcal{A}^{\text{dif}} = \mathbb{C}\langle \mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3 \dots \rangle$ denote the algebra of noncommutative polynomials on infinitely many variables $\mathbf{b}_n$, for $n \geq 1$, with unit 1. Then, the multiplicative map $\delta^{\text{dif}} : \mathcal{A}^{\text{dif}} \rightarrow \mathcal{H}^{\text{dif}} \otimes \mathcal{A}^{\text{dif}}$ defined on the generators by*

$$\delta^{\text{dif}} \mathbf{b}_n = 1 \otimes \mathbf{b}_n + \sum_{m=1}^{n-1} Q_m^{n-m-1}(\mathbf{a}_*) \otimes \mathbf{b}_{n-m},\tag{80}$$

where the polynomials $Q_m^l(\mathbf{a}_*)$ where defined in (70), gives a left coaction of $\mathcal{H}^{\text{dif}}$ on $\mathcal{A}^{\text{dif}}$.

Here are some examples of coactions:

$$\begin{aligned}\delta^{\text{dif}} \mathbf{b}_1 &= 1 \otimes \mathbf{b}_1, \\ \delta^{\text{dif}} \mathbf{b}_2 &= 1 \otimes \mathbf{b}_2 + \mathbf{a}_1 \otimes \mathbf{b}_1, \\ \delta^{\text{dif}} \mathbf{b}_3 &= 1 \otimes \mathbf{b}_3 + 2\mathbf{a}_1 \otimes \mathbf{b}_2 + \mathbf{a}_2 \otimes \mathbf{b}_1.\end{aligned}$$

Proof. We want to show that

$$(\text{Id} \otimes \delta^{\text{dif}}) \delta^{\text{dif}} = (\Delta^{\text{dif}} \otimes \text{Id}) \delta^{\text{dif}}.$$

On the left-hand side, we have

$$\begin{aligned}
(\text{Id} \otimes \delta^{\text{dif}}) \delta^{\text{dif}} \mathbf{b}_n &= 1 \otimes \delta^{\text{dif}} \mathbf{b}_n + \sum_{m=1}^{n-1} Q_m^{n-m-1}(\mathbf{a}_*) \otimes \delta^{\text{dif}} \mathbf{b}_{n-m} \\
&= 1 \otimes 1 \otimes \mathbf{b}_n + \sum_{m=1}^{n-1} 1 \otimes Q_m^{n-m-1}(\mathbf{a}_*) \otimes \mathbf{b}_{n-m} + \sum_{m=1}^{n-1} Q_m^{n-m-1}(\mathbf{a}_*) \otimes 1 \otimes \mathbf{b}_{n-m} \\
&\quad + \sum_{p=1}^{n-1} \sum_{q=1}^{n-p-1} Q_p^{n-p-1}(\mathbf{a}_*) \otimes Q_q^{n-p-q-1}(\mathbf{a}_*) \otimes \mathbf{b}_{n-p} \\
&= 1 \otimes 1 \otimes \mathbf{b}_n + \sum_{m=1}^{n-1} [1 \otimes Q_m^{n-m-1}(\mathbf{a}_*) + Q_m^{n-m-1}(\mathbf{a}_*) \otimes 1 \\
&\quad + \sum_{p=1}^{m-1} Q_p^{n-p-1}(\mathbf{a}_*) \otimes Q_{m-p}^{n-m-1}(\mathbf{a}_*)] \otimes \mathbf{b}_{n-m}.
\end{aligned}$$

Because of lemma (5.10), this term coincide with the term on the right-hand side, namely

$$(\Delta^{\text{dif}} \otimes \text{Id}) \delta^{\text{dif}} \mathbf{b}_n = 1 \otimes 1 \otimes \mathbf{b}_n + \sum_{m=1}^{n-1} \Delta^{\text{dif}} Q_m^{n-m-1}(\mathbf{a}_*) \otimes \mathbf{b}_{n-m}.$$

□

Remark 5.12 The abelianization $\mathcal{A}_{ab}^{\text{dif}} \cong \mathbb{C}[b_1, b_2, \dots]$ is the set of functions on the formal series

$$f(x) = 1 + \sum_n x^n b_n(f),$$

where the b_n are functions on the series defined as

$$b_n(f) = \frac{1}{n!} \frac{d^n f(0)}{dx^n}.$$

Then, the coaction of $\mathcal{H}_{ab}^{\text{dif}}$ on $\mathcal{A}_{ab}^{\text{dif}}$ induced by the coaction δ^{dif} of $\mathcal{H}^{\text{dif}}$ on $\mathcal{A}^{\text{dif}}$, is dual to the natural right action of $\mathcal{G}^{\text{dif}}$ on the set of series starting with 1.

In fact, if we start from two formal series $f(x) = 1 + \sum_n x^n b_n(f)$ and $g(x) = x + \sum_n x^{n+1} a_n(g)$ with commutative entries, the calculation of the composition is very similar to that of 5.5 and yields

$$\begin{aligned}
f(g(x)) &= 1 + \sum_{n=1}^{\infty} (g(x))^n b_n(f) \\
&= 1 + \sum_{n=1}^{\infty} x^n \left(a_n(f) + \sum_{m=1}^{n-1} Q_m^{n-m-1}(a_*(g)) b_{n-m}(f) \right) \\
&= 1 + \sum_{n=1}^{\infty} x^n \langle g \otimes f, 1 \otimes b_n + \sum_{m=1}^{n-1} Q_m^{n-m-1}(a_*) \otimes b_{n-m} \rangle.
\end{aligned}$$

5.3 Normal ordering

Given two matrix-valued series $\mathbf{f}(x) = x + \sum_{n=1}^{\infty} x^{n+1} a_n(\mathbf{f})$ and $\mathbf{g}(x) = x + \sum_{n=1}^{\infty} x^{n+1} a_n(\mathbf{g})$, we can not calculate $\mathbf{f} \circ \mathbf{g}(x)$ by replacing x by $\mathbf{g}(x)$ in the series defining $\mathbf{f}(x)$, before having fixed the position of the variable x in the series $\mathbf{f}(x)$. In fact, the problem is to extend the domain of definition of a function from a commutative domain to a non-commutative one. For instance, consider the function $\mathbf{f} : \mathbb{C} \longrightarrow M_4(\mathbb{C})$ defined as the multiplication by a non-diagonal matrix $A \in M_4(\mathbb{C})$, $\mathbf{f}(x) = xA = Ax$. When we try to extend it to the larger domain $M_4(\mathbb{C})$, which contains $\mathbb{C} \cong \mathbb{C} \cdot \mathbf{1}$, we observe a contradiction in the

evaluation on any matrix B : $\tilde{\mathbf{f}}(B) = BA \neq AB = \tilde{\mathbf{f}}(B)$. In other words, there are two possible ways to extend the function $\mathbf{f}$ and the normal ordering is just one of the two, chosen by fixing the position of the extended (non-commutative) variable. For instance, we can set $:\mathbf{f}(X): = XA$, for any matrix X .

Such a problem arises in QED when we want to consider the bare propagators, which depend on the bare electric charge e_0 , as functions of the renormalized electric charge e . In section 6 we show that the perturbative expression of $e_0 = e_0(e)$ depends on the noncommutative variables of the Hopf algebra $\mathcal{H}^\gamma$. In section 7 we will use the normal ordering to give a meaning to the composition $D(e_0) = D(e_0(e))$ in the noncommutative context.

Therefore, given any two matrix-valued formal series $\mathbf{f}(x)$ and $\mathbf{g}(x)$ starting with x , we define the normally ordered composition as

$$:\mathbf{f} \circ \mathbf{g}(x): = \mathbf{g}(x) \otimes 1 + \sum_{n=1}^{\infty} \mathbf{g}(x)^{n+1} \otimes a_n(\mathbf{f}) \quad (81)$$

$$= x + \sum_{n=1}^{\infty} x^{n+1} \langle \Delta^{\text{dif}} \mathbf{a}_n, \mathbf{g} \otimes \mathbf{f} \rangle. \quad (82)$$

Similarly, we can define the action of a series $\mathbf{g}(x)$ starting with x on a series $\mathbf{f}(x)$ starting with 1 as

$$\begin{aligned} :\mathbf{f} \circ \mathbf{g}(x): &= 1 + \sum_{n=1}^{\infty} (\mathbf{g}(x))^n \otimes b_n(\mathbf{f}) \\ &= 1 + \sum_{n=1}^{\infty} x^n \langle \delta^{\text{dif}} \mathbf{b}_n, \mathbf{g} \otimes \mathbf{f} \rangle \\ &= 1 + \sum_{n=1}^{\infty} x^n \langle \mathbf{g} \otimes \mathbf{f}, 1 \otimes \mathbf{b}_n + \sum_{m=1}^{n-1} Q_m^{n-m-1}(\mathbf{a}_*) \otimes \mathbf{b}_{n-m} \rangle. \end{aligned} \quad (83)$$

Remark 5.13 When the formal series converge to an analytical function, a non-perturbative expression for the normally ordered composition can be given as

$$:\mathbf{f} \circ \mathbf{g}(x): = \frac{1}{2\pi i} \int_{\gamma} \frac{dz}{z - \mathbf{g}(x)} \otimes \mathbf{f}(z),$$

where γ is a path in the complex plane surrounding the origin. This can be shown by adapting Kato's definition of the scalar functions on operators to matrix-valued functions $\mathbf{f}(z)$ (see [18]).

6 Renormalization of the electric charge

6.1 Renormalization of e_0

As we said in section 1, Dyson's formulas (12,13) for the renormalization of QED (plus the Ward identities) imply that the relation between the bare electric charge e_0 and the renormalized electric charge e is given only by means of the photon renormalization factor $Z_3(e)$, according to Eq. (11),

$$e_0 = Z_3(e)^{-1/2} e.$$

We show that such an equation makes sense in a noncommutative context.

Recall that $Z_3(e)$ is a formal power series on e with entries which depend on the elements of the noncommutative Hopf algebra $\mathcal{H}^\gamma$. More precisely, from Eqs.(13) and (66) we know that it is expanded over the photon generating trees $V(t) = \mathbf{1} \vee t$, so that

$$\begin{aligned} Z_3(e) &= 1 - \sum_{n=1}^{\infty} e^{2n} \zeta_3(\mathbf{v}_n) \\ &= \zeta_3(1 - \sum_{n=1}^{\infty} e^{2n} \mathbf{v}_n), \end{aligned} \quad (84)$$

where $\mathbf{v}_n = \sum_{|t|=n-1} V(t) \in \mathcal{H}^\gamma$ was defined in (43) and the $\mathbb{C}$ -valued function ζ_3 is extended linearly from $\mathcal{H}^\gamma$ to $\mathcal{H}^\gamma[[e]]$.

Proposition 6.1 *For any $n \geq 1$, define an element $\mathbf{w}_n$ in $\mathcal{H}^\gamma$ as*

$$\mathbf{w}_n = \sum_{k=1}^n \frac{(2k-1)!!}{k!2^k} P_n^{(k)}(\mathbf{v}_*), \quad (85)$$

and two elements $\mathbf{Z}_3(e)$ and $\mathbf{e}_0(e)$ in $\mathcal{H}^\gamma[[e^2]]$ as

$$\mathbf{Z}_3(e) = 1 - \sum_{n=1}^{\infty} e^{2n} \mathbf{v}_n, \quad (86)$$

$$\mathbf{e}_0(e) = e + \sum_{n=1}^{\infty} e^{2n+1} \mathbf{w}_n. \quad (87)$$

Then the Z_3 factor and the bare electric charge become

$$Z_3(e) = \zeta_3(\mathbf{Z}_3(e)), \quad (88)$$

$$e_0(e) = \zeta_3(\mathbf{e}_0(e)), \quad (89)$$

and the charge renormalization (11) can be deduced from a noncommutative analogue in $\mathcal{H}^\gamma[[e]]$,

$$\mathbf{e}_0(e) = e \mathbf{Z}_3(e)^{-1/2}. \quad (90)$$

Proof. Starting from the definitions (85), (86) and (87), we show that equation (90) holds.

$$\begin{aligned} \mathbf{Z}_3(e)^{-1/2} &= \left(1 - \sum_{p=1}^{\infty} e^{2p} \mathbf{v}_p\right)^{-1/2} \\ &= 1 + \sum_{k=1}^{\infty} \frac{(2k-1)!!}{k!2^k} \left(\sum_{p=1}^{\infty} e^{2p} \mathbf{v}_p\right)^k \\ &= 1 + \sum_{k=1}^{\infty} \frac{(2k-1)!!}{k!2^k} \sum_{p_1, \dots, p_k \geq 1} e^{2(p_1 + \dots + p_k)} \mathbf{v}_{p_1} \cdots \mathbf{v}_{p_k} \\ &= 1 + \sum_{k=1}^{\infty} \frac{(2k-1)!!}{k!2^k} \sum_{n \geq k} e^{2n} \sum_{p_1 + \dots + p_k = n} \mathbf{v}_{p_1} \cdots \mathbf{v}_{p_k} \\ &= 1 + \sum_{n=1}^{\infty} e^{2n} \sum_{k=1}^n \frac{(2k-1)!!}{k!2^k} P_n^{(k)}(\mathbf{v}_*) \\ &= 1 + \sum_{n=1}^{\infty} e^{2n} \mathbf{w}_n = \frac{\mathbf{e}_0(e)}{e}. \end{aligned}$$

□

Remark that inverting Eq. (85) we obtain

$$\mathbf{v}_n = \sum_{k=1}^n (-1)^{k+1} (k+1) P_n^{(k)}(\mathbf{w}_*).$$

Theorem 6.2 *The Connes-Kreimer map $\Psi : \mathcal{H}^{\text{odd}} \longrightarrow \mathcal{H}^\gamma$ given by*

$$\Psi(\mathbf{a}_{2n}) = \mathbf{w}_n$$

is an injective morphism of Hopf algebras. Therefore its image identifies the set of renormalization transformation of the electric charge as the noncommutative Hopf algebra of odd formal diffeomorphisms.

First we find a useful relation between the coefficients $\mathbf{u}_n$ and $\mathbf{w}_n$.

Lemma 6.3 *For any $n \geq 1$, the following relations hold:*

$$\mathbf{w}_n = \frac{1}{2}\mathbf{u}_n - \frac{1}{2} \sum_{p=1}^{n-1} \mathbf{w}_p \mathbf{w}_{n-p}, \quad (91)$$

$$\mathbf{u}_n = 2\mathbf{w}_n + \sum_{p=1}^{n-1} \mathbf{w}_p \mathbf{w}_{n-p}. \quad (92)$$

Proof. Relation (91) is the inverse of relation (92). Let us prove the second one.

If we introduce the definitions of $\mathbf{u}_n$ and $\mathbf{v}_n$ (Eqs.(42,43)) into Eq.(22), we obtain

$$\left(1 + \sum_{n=1}^{\infty} e^{2n} \mathbf{u}_n\right)^{-1} = 1 - \sum_{n=1}^{\infty} e^{2n} \mathbf{v}_n.$$

Thus

$$\frac{1}{\mathbf{Z}_3(e)} = 1 + \sum_{n=1}^{\infty} e^{2n} \mathbf{u}_n.$$

We use the fact

$$\frac{\mathbf{e}_0^2}{e^2} = \frac{1}{\mathbf{Z}_3(e)}$$

to have the equality

$$\begin{aligned} 1 + \sum_{n \geq 1} \mathbf{u}_n e^{2n} &= \left(1 + \sum_{n \geq 1} \mathbf{w}_n e^{2n}\right)^2 \\ &= 1 + \sum_{n \geq 1} \left[2\mathbf{w}_n + \sum_{p=1}^{n-1} \mathbf{w}_p \mathbf{w}_{n-p}\right] e^{2n}, \end{aligned}$$

from which follows relation (92). □

Proof of (6.2). We have to prove that

$$(\Psi \otimes \Psi) \Delta^{\text{dif}}(\mathbf{a}_{2n}) = \Delta^\gamma(\Psi(\mathbf{a}_{2n})),$$

that is,

$$\Delta^\gamma(\mathbf{w}_n) = \mathbf{w}_n \otimes 1 + 1 \otimes \mathbf{w}_n + \sum_{m=1}^{n-1} \bar{Q}_m^n(\mathbf{w}_*) \otimes \mathbf{w}_{n-m},$$

where the polynomials $\bar{Q}_m^n(\mathbf{w}_*) = Q_m^{2n-2m}(\mathbf{w}_*)$ were defined in (70). We prove it by induction, using the quadratic expression on the $\mathbf{w}$'s and $\mathbf{u}$'s of Eq. (91).

Suppose that the formula is true for all $\mathbf{w}_k$ with $k < n$. Write $\mathbf{w}_n$ as in (91), look at

$$\Delta^\gamma(\mathbf{w}_n) = \frac{1}{2} \Delta^\gamma(\mathbf{u}_n) - \frac{1}{2} \sum_{p=1}^{n-1} \Delta^\gamma(\mathbf{w}_p) \Delta^\gamma(\mathbf{w}_{n-p})$$

and compute each summand separately.

Take the term $\Delta^\gamma(\mathbf{u}_n)$ as computed in (4.2), it yields

$$\begin{aligned} \frac{1}{2}\Delta^\gamma \mathbf{u}_n &= \frac{1}{2}\mathbf{u}_n \otimes \mathbf{1} + \mathbf{1} \otimes \frac{1}{2}\mathbf{u}_n + \sum_{p=1}^{n-1} R_p^n(\mathbf{v}_*) \otimes \frac{1}{2}\mathbf{u}_{n-p} \\ &= \mathbf{w}_n \otimes \mathbf{1} + \frac{1}{2} \sum_{p=1}^{n-1} [\mathbf{w}_p \mathbf{w}_{n-p} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{w}_p \mathbf{w}_{n-p}] + \mathbf{1} \otimes \mathbf{w}_n \\ &\quad + \sum_{p=1}^{n-1} R_p^n(\mathbf{v}_*) \otimes \left[\mathbf{w}_{n-p} + \frac{1}{2} \sum_{k=1}^{n-p-1} \mathbf{w}_k \mathbf{w}_{n-p-k} \right]. \end{aligned}$$

The second term, by inductive hypothesis, is:

$$\begin{aligned} \frac{1}{2} \sum_{p=1}^{n-1} \Delta^\gamma(\mathbf{w}_p) \Delta^\gamma(\mathbf{w}_{n-p}) &= \\ &= \frac{1}{2} \sum_{p=1}^{n-1} \left[\mathbf{w}_p \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{w}_p + \frac{1}{2} \sum_{m=1}^{p-1} \bar{Q}_m^p \otimes \mathbf{w}_{p-m} \right] \left[\mathbf{w}_{n-p} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{w}_{n-p} + \frac{1}{2} \sum_{l=1}^{n-p-1} \bar{Q}_l^{n-p} \otimes \mathbf{w}_{n-p-l} \right]. \end{aligned}$$

Expanding the product gives

$$\begin{aligned} \frac{1}{2} \sum_{p=1}^{n-1} \Delta^\gamma(\mathbf{w}_p) \Delta^\gamma(\mathbf{w}_{n-p}) &= \\ &= \frac{1}{2} \sum_{p=1}^{n-1} [\mathbf{w}_p \mathbf{w}_{n-p} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{w}_p \mathbf{w}_{n-p}] + \sum_{p=1}^{n-1} \mathbf{w}_{n-p} \otimes \mathbf{w}_p \\ &\quad + \frac{1}{2} \sum_{p=1}^{n-1} \left[\sum_{m=1}^{p-1} \bar{Q}_m^p \mathbf{w}_{n-p} \otimes \mathbf{w}_{p-m} + \sum_{l=1}^{n-p-1} \mathbf{w}_p \bar{Q}_l^{n-p} \otimes \mathbf{w}_{n-p-l} \right] \\ &\quad + \frac{1}{2} \sum_{p=1}^{n-1} \sum_{m=1}^{p-1} \sum_{l=1}^{n-p-1} [\bar{Q}_m^p \otimes \mathbf{w}_{p-m} \mathbf{w}_{n-p} + \bar{Q}_l^{n-p} \otimes \mathbf{w}_p \mathbf{w}_{n-p-l} + \bar{Q}_m^p \bar{Q}_l^{n-p} \otimes \mathbf{w}_{p-m} \mathbf{w}_{n-p-l}]. \end{aligned}$$

Hence we obtain

$$\begin{aligned} \Delta^\gamma(\mathbf{w}_n) &= \mathbf{w}_n \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{w}_n \\ &\quad + \sum_{m=1}^{n-1} \left[R_m^n - \mathbf{w}_m - \frac{1}{2} \sum_{l=1}^{m-1} (\bar{Q}_{m-l}^{n-l} \mathbf{w}_l + \mathbf{w}_l \bar{Q}_{m-l}^{n-l}) \right] \otimes \mathbf{w}_{n-m} \\ &\quad + \frac{1}{2} \sum_{m=1}^{n-1} \sum_{p=1}^{n-m-1} \left[R_m^n - \bar{Q}_m^{m+p} - \bar{Q}_m^{n-p} - \sum_{l=1}^{m-1} \bar{Q}_l^{l+p} \bar{Q}_{m-l}^{n-p-l} \right] \otimes \mathbf{w}_p \mathbf{w}_{n-m-p} \end{aligned}$$

and therefore

$$\Delta^\gamma(\mathbf{w}_n) = \mathbf{w}_n \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{w}_n + \sum_{m=1}^{n-1} \bar{Q}_m^{n-m} \otimes \mathbf{w}_{n-m}$$

because of the two relations between the R 's and the $\bar{Q}$'s proved in the following lemmas (6.5) and (6.6).

□

Lemma 6.4 *The generating function*

$$\bar{Q}(x, y) = \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} x^n y^{2m} \bar{Q}_m^{n+m}(\mathbf{w}_*) + \frac{1}{1-x} \quad (93)$$

satisfies the equation

$$\bar{Q}(x, y) = (1 - x\mathbf{Z}_0(y)^2)^{-1}\mathbf{Z}_0(y), \quad (94)$$

where $\mathbf{Z}_0(y) = 1 + \sum_{n=1}^{\infty} y^{2n} \mathbf{w}_n \in \mathcal{H}^\gamma[[y^2]]$.

Proof. For the proof, we use the combinatorial identity

$$\binom{2n-2m+1}{k} = \binom{2n-2m-1}{k} + 2\binom{2n-2m-1}{k-1} + \binom{2n-2m-1}{k-2}$$

to derive the recurrence relation

$$\bar{Q}_m^n = \bar{Q}_m^{n-1} + 2\mathbf{w}_m + P_m^{(2)}(\mathbf{w}_*) + 2 \sum_{k=1}^{m-1} \mathbf{w}_k \bar{Q}_{m-k}^{n-k-1} + \sum_{k=2}^{m-2} P_k^{(2)}(\mathbf{w}_*) \bar{Q}_{m-k}^{n-k-1}.$$

The expression for $\bar{Q}(x, y)$ follows by the usual method: in the last equation, we write $n = m + \alpha$, we multiply by $x^\alpha y^{2m}$ and we sum over α and m from 1 to ∞ . This gives us

$$\begin{aligned} \bar{Q}(x, y) - \frac{1}{1-x} - \mathbf{Z}_0(y) + 1 &= \sum_{m=1}^{\infty} \sum_{\alpha=1}^{\infty} x^\alpha y^{2m} \bar{Q}_m^{m+\alpha} \\ &= x \left(\bar{Q}(x, y) - \frac{1}{1-x} \right) + \frac{2x(\mathbf{Z}_0(y) - 1)}{1-x} + \frac{x(\mathbf{Z}_0(y) - 1)^2}{1-x} \\ &\quad + 2x(\mathbf{Z}_0(y) - 1) \left(\bar{Q}(x, y) - \frac{1}{1-x} \right) + x(\mathbf{Z}_0(y) - 1)^2 \left(\bar{Q}(x, y) - \frac{1}{1-x} \right) \\ &= 1 - \frac{1}{1-x} + x(\mathbf{Z}_0(y))^2 \bar{Q}(x, y). \end{aligned}$$

Solving for $\bar{Q}(x, y)$ yields Eq.(94). $\square$

Lemma 6.5 *The following identity holds for any $n, m \geq 1$:*

$$R_m^n(\mathbf{v}_*) = \bar{Q}_m^n(\mathbf{w}_*) + \mathbf{w}_m + \frac{1}{2} \sum_{l=1}^{m-1} (\bar{Q}_{m-l}^{n-l}(\mathbf{w}_*) \mathbf{w}_l + \mathbf{w}_l \bar{Q}_{m-l}^{n-l}(\mathbf{w}_*)).$$

Proof. Using Eq.(94) and $\mathbf{Z}_0(y) = (\mathbf{Z}_3(y))^{-1/2}$ we can write

$$\bar{Q}(x, y) = \left(1 - \frac{x}{\mathbf{Z}_3(y)} \right) (\mathbf{Z}_3(y))^{-1/2} = (\mathbf{Z}_3(y) - x)^{-1} (\mathbf{Z}_3(y))^{1/2} = R(x, y) (\mathbf{Z}_3(y))^{1/2}.$$

Therefore, $R(x, y) = \bar{Q}(x, y) \mathbf{Z}_0(y)$. Since $\mathbf{Z}_0(y)$ commutes with $\bar{Q}(x, y)$ (because of Eq.(94)), we obtain

$$R(x, y) = \frac{1}{2} (\bar{Q}(x, y) \mathbf{Z}_0(y) + \mathbf{Z}_0(y) \bar{Q}(x, y)).$$

In the expansion of this equality over x and y , the term $x^n y^{2m}$, for any $n \geq 0$ and $m \geq 1$, gives the equality

$$R_m^{n+m} = \bar{Q}_m^{n+m} + \mathbf{w}_m + \frac{1}{2} \sum_{l=1}^{m-1} (\bar{Q}_{m-l}^{n+m-l} \mathbf{w}_l + \mathbf{w}_l \bar{Q}_{m-l}^{n+m-l})$$

from which the lemma follows. $\square$

Lemma 6.6 *The following identity holds for any $n \geq 0$ and $m \geq 1$:*

$$R_m^n(\mathbf{v}_*) = \bar{Q}_m^{n+p}(\mathbf{w}_*) + \bar{Q}_m^{n-p}(\mathbf{w}_*) + \sum_{l=1}^{m-1} \bar{Q}_l^{p+l}(\mathbf{w}_*) \bar{Q}_{m-l}^{n-p-l}(\mathbf{w}_*).$$

Proof. The definition of the generating function enables us to write

$$\bar{Q}(x, y)\bar{Q}(z, y) - \frac{1}{(1-x)(1-z)} = \sum_{m=1}^{\infty} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} x^p y^{2m} z^q \left(\bar{Q}_m^{m+p} + \bar{Q}_m^{m+q} + \sum_{l=1}^{m-1} \bar{Q}_l^{l+p} \bar{Q}_{m-l}^{m-l+q} \right).$$

On the other hand, from the relation between $\bar{Q}(x, y)$ and $R(x, y)$ we find

$$\bar{Q}(x, y)\bar{Q}(z, y) = \mathbf{Z}_3(y)R(x, y)R(z, y).$$

Now we use Eq.(55), the resolvent identity for $R(x, y)R(z, y)$, to write

$$\mathbf{Z}_3(y)R(x, y)R(z, y) = \mathbf{Z}_3(y) \frac{R(x, y) - R(z, y)}{x - z} = \frac{xR(x, y) - zR(z, y)}{x - z},$$

where we have used the relations $\mathbf{Z}_3(y)R(x, y) = 1 + xR(x, y)$. Finally, we obtain

$$\begin{aligned} \bar{Q}(x, y)\bar{Q}(z, y) - \frac{1}{(1-x)(1-z)} &= \frac{xR(x, y) - zR(z, y)}{x - z} - \frac{1}{(1-x)(1-z)} \\ &= \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} y^{2m} \frac{x^{k+1} - z^{k+1}}{x - z} R_m^{m+k} \\ &= \sum_{m=1}^{\infty} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} y^{2m} x^p z^q R_m^{m+p+q}, \end{aligned}$$

since

$$\frac{x^{k+1} - z^{k+1}}{x - z} = \sum_{p+q=k} x^p z^q.$$

If we identify the coefficient of $x^p y^{2m} z^q$ on both sides, we obtain the identity

$$R_m^{m+p+q}(\mathbf{v}_*) = \bar{Q}_m^{m+p}(\mathbf{w}_*) + \bar{Q}_m^{m+q}(\mathbf{w}_*) + \sum_{l=1}^{m-1} \bar{Q}_l^{l+p}(\mathbf{w}_*) \bar{Q}_{m-l}^{m-l+q}(\mathbf{w}_*).$$

from which follows the lemma. $\square$

6.2 Renormalization of α_0

The Connes-Kreimer morphism Ψ between the Hopf algebras $\mathcal{H}^{\text{odd}}$ and $\mathcal{H}^\gamma$ was chosen according to the prescription $\mathbf{e}_0 = e\mathbf{Z}_3(e)^{-1/2}$, which corresponds to the choice of Ref.[12]. However, a simpler morphism can be given between the Hopf algebras $\mathcal{H}^{\text{dif}}$ and $\mathcal{H}^\gamma$. This is the subject of the present section.

In QED, the propagators and renormalization factors depend on the squares e_0^2 and e^2 of the coupling constants. Therefore, it is tempting to expand them over the fine structure constants $\alpha_0 = e_0^2/4\pi$ and $\alpha = e^2/4\pi$.

Let us set up the notations with respect to the new variable α :

$$\begin{aligned} \mathbf{Z}_3(\alpha) &= 1 - \sum_{n=1}^{\infty} (4\pi)^n \alpha^n \mathbf{v}_n, \\ \frac{1}{\mathbf{Z}_3(\alpha)} &= 1 + \sum_{n=1}^{\infty} (4\pi)^n \alpha^n \mathbf{u}_n, \\ \alpha_0 = \frac{\alpha}{\mathbf{Z}_3(\alpha)} &= \alpha + \sum_{n=1}^{\infty} (4\pi)^n \alpha^{n+1} \mathbf{u}_n. \end{aligned} \tag{95}$$

Theorem 6.7 *Let us define the Connes-Kreimer map $\hat{\Psi} : \mathcal{H}^{\text{dif}} \longrightarrow \mathcal{H}^\gamma$ as the algebra morphism given on the generators by $\hat{\Psi}(\mathbf{a}_n) = (4\pi)^n \mathbf{u}_n$ and $\hat{\Psi}(1) = 1$. Then $\hat{\Psi}$ is an injective Hopf algebra morphism, and the Hopf algebra $\mathcal{H}^{\text{dif}}$ can be seen as the set of renormalization transformations of the fine structure constant.*

Lemma 6.8 *For any $n \geq 0$ and any $m > 0$ we have an identity*

$$R_m^n(\mathbf{v}_*) = Q_m^{n-m}(\mathbf{u}_*).$$

Proof. To obtain this result, we start from the generating function $R(x, y)$, defined by Eq. (56), that we rewrite as

$$R(x, y) = (\mathbf{Z}_3(y) - x)^{-1} = \mathbf{Z}_3(y)^{-1} [1 - x\mathbf{Z}_3(y)^{-1}]^{-1}.$$

From lemma (2.9), by taking the sum over all trees in Eq. (25), we see that the inverse of the series $\mathbf{Z}_3(y) = 1 + \sum_{n=1}^{\infty} y^{2n} \mathbf{v}_n$ is

$$\frac{1}{\mathbf{Z}_3(y)} = 1 - \sum_{n=1}^{\infty} y^{2n} \mathbf{u}_n.$$

Now, in lemma (5.8) we show that if we set $\mathbf{f}(y) = y + \sum_{n=1}^{\infty} y^{2n+1} \mathbf{u}_n = y\mathbf{Z}_3(y)^{-1}$, then $Q(x, y)$ satisfies the equation

$$\begin{aligned} Q(x, y) &= \left(\frac{y}{\mathbf{f}(y)} - x \right)^{-1} = \mathbf{f}(y) [y - x\mathbf{f}(y)]^{-1} \\ &= \mathbf{Z}_3(y)^{-1} [1 - x\mathbf{Z}_3(y)^{-1}]^{-1} = R(x, y). \end{aligned}$$

If we expand the generating functions $Q(x, y)$ and $R(x, y)$ over x and y and consider the coefficient of $x^n y^{2m}$ we obtain the expected result. $\square$

Proof of (6.7). The map $\hat{\Psi}$ is obviously compatible with the coproduct, because of lemma (6.8) and proposition (4.2). In fact,

$$\begin{aligned} (\hat{\Psi} \otimes \hat{\Psi}) \circ \Delta^{\text{dif}} \mathbf{a}_n &= \hat{\Psi}(\mathbf{a}_n) \otimes 1 + 1 \otimes \hat{\Psi}(\mathbf{a}_n) + \sum_{m=1}^{n-1} Q_m^{n-m}(\hat{\Psi}(\mathbf{a}_*)) \otimes \hat{\Psi}(\mathbf{a}_{n-m}) \\ &= (4\pi)^n \mathbf{u}_n \otimes 1 + 1 \otimes (4\pi)^n \mathbf{u}_n + \sum_{m=1}^{n-1} Q_m^{n-m}((4\pi)^* \mathbf{u}_*) \otimes (4\pi)^{n-m} \mathbf{u}_{n-m} \\ &= \Delta^\gamma((4\pi)^n \mathbf{u}_n) = \Delta^\gamma(\hat{\Psi}(\mathbf{a}_n)). \end{aligned}$$

$\square$

7 Noncommutative Dyson's formulas

In this section, we show that Dyson's equations (9,10) can be seen as a consequence of the Connes-Kreimer map of section 5, between the Hopf algebra of formal diffeomorphisms and the Hopf algebra of QED renormalization.

But we shall prove more: Dyson's formulas are still valid when $Z_2(e)$ and $Z_3(e)$ are matrices, that is, for their noncommutative versions $\mathbf{Z}_2(e)$ and $\mathbf{Z}_3(e)$. In fact, scalar Dyson's formulas can be deduced from analogue identities in the algebra $\mathcal{H}^{\text{qed}} \otimes \mathcal{H}^{\text{qed}}$. This is the essence of noncommutative renormalization.

Dyson's formulas involve the bare and renormalized propagators, which are functions respectively of the bare and of the renormalized electric charge. In order to confront them, we are forced to describe the explicit dependency of the bare propagators on the renormalized charge, that is, we are forced to perform a composition between the bare propagators, as functions of e_0 , and e_0 as a function of e .

In section 6 we showed that the bare electric charge is the evaluation of a multiplicative scalar map ζ_3 on a non-commutative formal series,

$$\mathbf{e}_0(e) = e - \sum_{n=1}^{\infty} e^{2n+1} \mathbf{w}_n \in \mathcal{H}^\gamma[[e]].$$

Wishing to isolate some non-commutative analogue of the bare propagators, we are forced to regard the series in $\mathcal{H}^{\text{qed}}[[e_0]]$ as series in $\mathcal{H}^{\text{qed}}[[\mathbf{e}_0]]$, and then compose them with $\mathbf{e}_0(e)$ as a series in $\mathcal{H}^{\text{qed}}[[e]]$. As remarked in (82), when we want to extend the domain of definition of a series, from the set of complex numbers to a non-commutative set, the normal ordering is required to fix the position of the variables. Therefore, in this section we make use of the notation of section 5.3.

7.1 Dyson's formula for photons

Proposition 7.1 *The operators*

$$:\mathbf{D}(\mathbf{e}_0): = \mathbf{1} \otimes \mathbf{1} + \sum_{n=1}^{\infty} \mathbf{e}_0^{2n} \otimes \mathbf{u}_n \in \mathcal{H}^\gamma[[\mathbf{e}_0^2]], \quad (96)$$

$$\bar{\mathbf{D}}(e) = \mathbf{1} \otimes \mathbf{1} + \sum_{n=1}^{\infty} e^{2n} \Delta^\gamma \mathbf{u}_n \in \mathcal{H}^\gamma \otimes \mathcal{H}^\gamma[[e^2]] \quad (97)$$

satisfy the noncommutative Dyson formula

$$(\mathbf{Z}_3(e) \otimes \mathbf{1}) \bar{\mathbf{D}}(e) = :\mathbf{D}(\mathbf{e}_0): \quad (98)$$

for $\mathbf{e}_0 = e \mathbf{Z}_3(e)^{-1/2}$.

Moreover, the bare and renormalized photon propagators can be recovered as

$$\begin{aligned} D(e_0) &= \langle \zeta_3 \otimes \varphi^\gamma, :\mathbf{D}(\mathbf{e}_0): \rangle, \\ \bar{D}(e) &= \langle \zeta_3 \otimes \varphi^\gamma, \bar{\mathbf{D}}(e) \rangle, \end{aligned}$$

and Dyson's formula (10) follows from Eq. (98).

Proof. Equation (98) is quite easy to prove. Since

$$\Delta^\gamma \mathbf{u}_n = \mathbf{u}_n \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} R_m^n(\mathbf{v}_*) \otimes \mathbf{u}_{n-m},$$

if we expand the left hand-side of (98) we obtain

$$\begin{aligned} (\mathbf{Z}_3(e) \otimes \mathbf{1}) \bar{\mathbf{D}}(e) &= \mathbf{1} \otimes \mathbf{1} + \sum_{n=1}^{\infty} e^{2n} \left[(\mathbf{u}_n - \mathbf{v}_n - \sum_{k=1}^{n-1} \mathbf{v}_k \mathbf{u}_{n-k}) \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{u}_n \right. \\ &\quad \left. + \sum_{m=1}^{n-1} (R_m^n - \mathbf{v}_m - \sum_{k=1}^{m-1} \mathbf{v}_k R_{m-k}^n) \otimes \mathbf{u}_{n-m} \right]. \end{aligned}$$

The factor of $\otimes \mathbf{1}$ is zero because of Eq.(50), and the factor of $\otimes \mathbf{u}_{n-m}$ is equal to R_m^{n-1} because of Eq.(51). Hence,

$$\begin{aligned} (\mathbf{Z}_3(e) \otimes \mathbf{1}) \bar{\mathbf{D}}(e) &= \mathbf{1} \otimes \mathbf{1} + \sum_{n=1}^{\infty} e^{2n} \left(\mathbf{1} \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} R_m^{n-1} \otimes \mathbf{u}_{n-m} \right) \\ &= \mathbf{1} \otimes \mathbf{1} + \sum_{n=1}^{\infty} e^{2n} \left(\mathbf{1} \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} Q_m^{n-m-1}(\mathbf{u}_*) \otimes \mathbf{u}_{n-m} \right). \end{aligned}$$

On the right hand-side of (98), apply the normally ordered composition. Because the series $: \mathbf{D}(\mathbf{e}_0) :$ starts with 1, we apply the action (83). Since the Connes-Kreimer map transforms the coproduct on series into the photon coproduct, and because $: \mathbf{D}(\mathbf{e}_0) :$ depends on the square of $\mathbf{e}_0$, where

$$\mathbf{e}_0^2 = \frac{e^2}{\mathbf{Z}_3(e)} = e^2 + \sum_{n=1}^{\infty} (e^2)^{n+1} \mathbf{u}_n,$$

the series $: \mathbf{D}(\mathbf{e}_0) :$ is

$$: \mathbf{D}(\mathbf{e}_0) : = 1 \otimes 1 + \sum_{n=1}^{\infty} e^{2n} (1 \otimes \mathbf{u}_n + \sum_{m=1}^{n-1} Q_m^{n-m-1}(\mathbf{u}_*) \otimes \mathbf{u}_{n-m})$$

because the Connes-Kreimer map is a morphism of Hopf algebras. Therefore, modulo the free action μ , we obtain

$$(\mathbf{Z}_3(e) \otimes 1) \bar{\mathbf{D}}(e) = : \mathbf{D}(\mathbf{e}_0) :.$$

To recover the classical Dyson's formula, recall that the bare and renormalized photon propagators were expanded in section 1 as

$$\begin{aligned} D(q, e_0) &= 1 + \sum_{n=1}^{\infty} e_0^{2n} D_n(q), \\ \bar{D}(q, e_0) &= 1 + \sum_{n=1}^{\infty} e_0^{2n} \bar{D}_n(q), \end{aligned}$$

and that the results of section 4 tell us how to find the coefficients:

$$\begin{aligned} D_n(q) &= \varphi^\gamma(\mathbf{u}_n) = \langle \varphi^\gamma, \mathbf{u}_n \rangle, \\ \bar{D}_n(q) &= \bar{\varphi}^\gamma(\mathbf{u}_n) = \langle \zeta_3 \otimes \varphi^\gamma, \Delta^\gamma \mathbf{u}_n \rangle. \end{aligned}$$

□

Since Dyson's formula is noncommutative, we can use operators and not only scalars as Z_3 . An example is $Z_3 = 1 + \Pi(q)$. Dyson's formula becomes now

$$(1 + \Pi(q)) \bar{D}(q; e) = : D(q; \frac{e}{\sqrt{1 + \Pi(q)}}) :.$$

Similar algebraic relations can be established for the vacuum polarization.

Proposition 7.2 *The operators*

$$: \Pi(\mathbf{e}_0) : = \sum_{n=1}^{\infty} \mathbf{e}_0^{2n} \otimes \mathbf{v}_n \in \mathcal{H}^\gamma[[\mathbf{e}_0^2]], \quad (99)$$

$$\bar{\Pi}(e) = \sum_{n=1}^{\infty} e^{2n} \Delta^\gamma \mathbf{v}_n \in \mathcal{H}^\gamma \otimes \mathcal{H}^\gamma[[e^2]] \quad (100)$$

satisfy the noncommutative Dyson formula

$$1 \otimes 1 - \bar{\Pi}(e) = (\mathbf{Z}_3(e) \otimes 1)(1 \otimes 1 - : \Pi(\mathbf{e}_0) :), \quad (101)$$

for $\mathbf{e}_0 = e \mathbf{Z}_3(e)^{-1/2}$.

The bare and renormalized vacuum polarizations can be recovered as

$$\begin{aligned} \Pi(e_0) &= \langle \zeta_3 \otimes \varphi^\gamma, : \Pi(\mathbf{e}_0) : \rangle, \\ \bar{\Pi}(e) &= \langle \zeta_3 \otimes \varphi^\gamma, \bar{\Pi}(e) \rangle. \end{aligned}$$

Proof. The proof is similar to that of (7.1).

$$:\mathbf{\Pi}(e_0): = \sum_{n=1}^{\infty} e_0^{2n} \otimes \mathbf{v}_n \quad (102)$$

$$= \sum_{n=1}^{\infty} e^{2n} (\mathbf{1} \otimes \mathbf{v}_n + \sum_{m=1}^{n-1} Q_m^{n-m-1}(\mathbf{u}_*) \otimes \mathbf{v}_{n-m}) \quad (103)$$

$$= \sum_{n=1}^{\infty} e^{2n} (\mathbf{1} \otimes \mathbf{v}_n + \sum_{m=1}^{n-1} R_m^{n-1} \otimes \mathbf{v}_{n-m}) \quad (104)$$

$$\bar{\mathbf{\Pi}}(e) = \sum_{n=1}^{\infty} e^{2n} [\mathbf{v}_n \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{v}_n + \sum_{m=1}^{n-2} R_m^{n-2}(\mathbf{v}_*) \otimes \mathbf{v}_{n-m}]. \quad (105)$$

Then, using Eq.(51), it is straightforward to show that

$$\mathbf{1} \otimes \mathbf{1} - \bar{\mathbf{\Pi}}(e) = (\mathbf{Z}_3(e) \otimes \mathbf{1})(\mathbf{1} \otimes \mathbf{1} - \mathbf{\Pi}(e_0)).$$

□

As a noncommutative example of this second type of Dyson's formula, we take $Z_3 = 1 - \Pi(p; e)$:

$$1 - \bar{\Pi}(p, q; e) = (1 - \Pi(p; e)) \left(1 - \Pi(q; \frac{e}{\sqrt{1 - \Pi(p; e)}}) \right).$$

7.2 Dyson's formula for electrons

The case of the electron is very similar. We give only the definitions and the results.

Proposition 7.3 *The operators*

$$:\mathbf{S}(\mathbf{e}_0): = \mathbf{1} \otimes \mathbf{1} + \sum_{n=1}^{\infty} \mathbf{e}_0^{2n} \otimes \mathbf{x}_n \in \mathcal{H}^e[[\mathbf{e}_0^2]], \quad (106)$$

$$\bar{\mathbf{S}}(e) = \mathbf{1} \otimes \mathbf{1} + \sum_{n=1}^{\infty} e^{2n} \Delta^e \mathbf{x}_n \in \mathcal{H}^{\text{qed}} \otimes \mathcal{H}^e[[e^2]] \quad (107)$$

$$\mathbf{Z}_2(e) = \mathbf{1} + \sum_{n=1}^{\infty} e^{2n} \mathbf{y}_n \in \mathcal{H}^e[[e^2]] \quad (108)$$

satisfy the noncommutative Dyson formula in $\mathcal{H}^{\text{qed}} \otimes \mathcal{H}^e[[e^2]]$

$$(\mathbf{Z}_2(e) \otimes \mathbf{1}) \bar{\mathbf{S}}(e) = : \mathbf{S}(\mathbf{e}_0) : \quad (109)$$

for $\mathbf{e}_0 = e \mathbf{Z}_3(e)^{-1/2}$.

Moreover, the bare and renormalized electron propagators can be recovered as

$$\begin{aligned} S(e_0) &= \langle (\zeta_3, \zeta_2) \otimes \varphi^e, : \mathbf{S}(\mathbf{e}_0) : \rangle, \\ \bar{S}(e) &= \langle (\zeta_3, \zeta_2) \otimes \varphi^e, \bar{\mathbf{S}}(e) \rangle, \end{aligned}$$

and Dyson's formula (9) follows from Eq. (109).

Proposition 7.4 *The operators*

$$:\mathbf{\Sigma}(\mathbf{e}_0): = - \sum_{n=1}^{\infty} \mathbf{e}_0^{2n} \otimes \mathbf{y}_n \in \mathcal{H}^e[[\mathbf{e}_0^2]], \quad (110)$$

$$\bar{\mathbf{\Sigma}}(e) = - \sum_{n=1}^{\infty} e^{2n} \Delta^e \mathbf{y}_n \in \mathcal{H}^{\text{qed}} \otimes \mathcal{H}^e[[e^2]] \quad (111)$$

satisfy the noncommutative Dyson formula in $\mathcal{H}^{\text{qed}} \otimes \mathcal{H}^e[[e^2]]$

$$1 \otimes 1 - \bar{\Sigma}(e) = (\mathbf{Z}_2(e) \otimes 1)(1 \otimes 1 - : \Sigma(\mathbf{e}_0) :).$$
 (112)

The bare and renormalized electron self-energies can be recovered as

$$\begin{aligned} \Sigma(e_0) &= \langle (\zeta_3, \zeta_2) \otimes \varphi^e, : \Sigma(\mathbf{e}_0) : \rangle, \\ \bar{\Sigma}(e) &= \langle (\zeta_3, \zeta_2) \otimes \varphi^e, \bar{\Sigma}(e) \rangle. \end{aligned}$$

8 Conclusion and perspective

In this paper we provided explicit formulas, valid at all orders, for the renormalization of massless QED. These formulas are equivalent to Zimmermann's forest formula for the propagators, self-energy or vacuum polarization but they use the coefficients of e^{2n} in the perturbative expansion instead of the Feynman diagrams. The coefficients of e^{2n} make full use of Ward's identity $Z_1 = Z_2$ and decrease the number of integrals to calculate. This number grows as a polynomial in n instead of an exponential of n as is the case for Feynman diagrams.

Besides this practical aspect, this paper presented a noncommutative extension of renormalization which enables us to consider the three lepton generations as a single quantity and to renormalize them in one stroke. This provides a proper framework to investigate lepton transmutation [3]. The Hopf algebra structure of noncommutative renormalization preserves the main properties of renormalization, such as its coassociativity, which is the root of the renormalization group. The Connes-Kreimer map still exists, but the objects corresponding to the group of diffeomorphisms remains to be explored in the noncommutative setting.

In forthcoming publications, we shall present two extensions of the present work. Firstly, we shall consider the case of massive QED. The formulas for renormalized massive propagators are simple extensions of those for massless propagators, involving the derivatives of the bare propagators with respect to the mass m . However, the extension of the Hopf algebra of renormalization is more involved. Secondly, we shall investigate more fully the bigrading of QED, where photon and electron loops are distinguished. In particular, we shall give explicit formulas for the renormalization at a fixed number of photon and electron loops. The loop bigrading is interesting because the series over electron loops can be summed to all orders, providing a systematic way to obtain semi-perturbative (or semi-nonperturbative) equations for the propagators, where the electron loops are summed to all orders and the photon loops are introduced one by one.

Acknowledgements

We are grateful to the Institut Girard Desargues of the University of Lyon I for kind hospitality and support. We thank Dirk Kreimer and Jean-Louis Loday for helpful discussions. This is IPGP contribution #0000.

References

- [1] E. Bagan, M. Lavelle, and D. McMullan. Charges from dressed matter: Construction. *Ann. Phys.*, 282:471–502, 2000.
- [2] E. Bagan, M. Lavelle and D. McMullan. Charges from dressed matter: Physics and renormalisation. *Ann. Phys.*, 282:503–540, 2000.
- [3] J. Bordes, H.M. Chan, J. Faridani and S.T. Tsou. Photo-Transmutation of Leptons hep-ph/0007004
- [4] D.J. Broadhurst, R. Delbourgo, and D. Kreimer. Unknotting the polarized vacuum of quenched QED. *Phys. Lett. B*, 366:421–430, 1999.
- [5] Ch. Brouder. Runge-Kutta methods and renormalization. *Eur. Phys. J. C*, 12:521–34, 2000.

- [6] Ch. Brouder. On the trees of quantum fields. *Eur. Phys. J. C*, 12:535–49, 2000.
- [7] Ch. Brouder and A. Frabetti. Renormalization of QED with planar binary trees. *Eur. Phys. J. C*, to be published.
- [8] J.C. Collins. *Renormalization*. Cambridge University Press, Cambridge, 1984.
- [9] A. Connes and H. Moscovici. Hopf algebras, cyclic cohomology and the transverse index theorem. *Commun. Math. Phys.*, 198:199–246, 1998.
- [10] A. Connes and D. Kreimer. Hopf algebras, renormalization and noncommutative geometry. *Commun. Math. Phys.*, 199:203–42, 1998.
- [11] A. Connes and D. Kreimer. Renormalization in quantum field theory and the Riemann-Hilbert problem I: the Hopf algebra structure of graphs and the main theorem. *Commun. Math. Phys.*, 210:249–273, 2000.
- [12] A. Connes and D. Kreimer. Renormalization in quantum field theory and the Riemann-Hilbert problem II: the β function, diffeomorphisms and the renormalization group. *Commun. Math. Phys.*, to be published. hep-th/0003188.
- [13] F.J. Dyson. The S matrix in quantum electrodynamics. *Phys. Rev.*, 75:1736–55, 1949.
- [14] R.P. Feynman. Space-time approach to quantum electrodynamics. *Phys. Rev.*, 76:p.769–89, 1949.
- [15] A. Frabetti. Simplicial properties of the set of planar binary trees. *J. Alg. Comb.*, to appear.
- [16] V. Ginzburg and M.M. Kapranov. Koszul duality for operads. *Duke Math. J.*, 76:p.203–272, 1994.
- [17] C. Itzykson and J.-B. Zuber. *Quantum Field Theory*. McGraw-Hill, New York, 1980.
- [18] T. Kato. *Perturbation Theory for Linear Operators*. Springer, Berlin, 1980.
- [19] D. Kreimer. On the Hopf algebra structure of perturbative quantum field theory. *Adv. Th. Math. Phys.*, 2:303–34, 1998.
- [20] D. Kreimer. Chen’s iterated integral represents the operator product expansion. *Adv. Th. Math. Phys.*, 3, 1999. to be published (hep-th/9901099).
- [21] D. Kreimer. *Knots and Feynman diagrams*. Cambridge University Press, 2000.
- [22] J.-L. Loday. Arithmetree. Preprint:<http://www-irma.u-strasbg.fr/~loday/arithmetree.ps>.
- [23] J.-L. Loday. Algèbres ayant deux opérations associatives (digèbres). *C. R. Acad. Sci. Paris*, 321:p.141–146, 1995.
- [24] J.-L. Loday and T. Pirashvili. Universal Enveloping Algebras of Leibniz Algebras and (Co)homology. *Math. Annalen*, 296:569–572, 1993.
- [25] J.-L. Loday and M.O. Ronco. Hopf algebra of the planar binary trees. *Adv. Math.*, 139:293–309, 1998.
- [26] J.-L. Loday and M.O. Ronco. Order structure and the algebra of permutations and planar binary trees. Preprint:<http://www-irma.u-strasbg.fr/~loday/LR2.ps>.
- [27] M.E. Peskin and D.V. Schroeder. *An Introduction to Quantum Field Theory*. Perseus Books Pub. L.L.C., 1995.
- [28] S.S. Schweber. *An Introduction to Relativistic Quantum Field Theory*. Harper and Row, New York, 1961.
- [29] M.E. Sweedler. *Hopf algebras*. W.A. Benjamin Inc., NY, 1969.

- [30] J.C. Ward. An identity in quantum electrodynamics. *Phys. Rev.*, 78:182, 1950.
- [31] S. Weinberg. *The Quantum Theory of Fields*. Cambridge Univ. Press, Cambridge, 1995.
- [32] W. Zimmermann. Convergence of Bogoliubov's method of renormalization in momentum space. *Commun. Math. Phys.*, 15:208–34, 1969.